

CROSSED-PRODUCT C*-ALGEBRAS FOR CONFORMAL AUTOMORPHISMS OF THE DISK

MAN-DUEN CHOI AND FRÉDÉRIC LATRÉMOLIÈRE

ABSTRACT. We study the C*-algebra crossed-product of the closed unit disk by the action of one of its conformal automorphisms. After classifying the conformal automorphisms up to topological conjugacy, we investigate, for each class, the irreducible representations of the full C*-crossed-products, and derive their spectrum and a complete description of the algebras.

1. INTRODUCTION

We present in this paper a detailed description of the C*-crossed product algebras associated to conformal discrete dynamical systems of the closed unit disk $\mathbb{D}$.

Given a compact Hausdorff space X and an homeomorphism φ on X , the C*-crossed product of $C(X)$ by $\mathbb{Z}$ generated by the *-automorphism $f \in C(X) \mapsto f \circ \varphi$ is the universal C*-algebra [18] [12] [15] generated by $C(X)$ and a unitary U_φ such that for all $f \in C(X)$ we have $U_\varphi^* f U_\varphi = f \circ \varphi$. We refer to U_φ as the canonical unitary of $C(X) \rtimes_\varphi \mathbb{Z}$. The universal property means that for any C*-algebra $\mathfrak{A}$ containing a unitary V and a C*-algebra $\mathcal{A}$ such that there exists a *-homomorphism $\rho : C(X) \longrightarrow \mathcal{A}$ which satisfies $\rho(U_\varphi^* f U_\varphi) = V^* \rho(f) V$ for all $f \in C(X)$, there exists a (necessarily unique) *-homomorphism $\psi : C(X) \rtimes_\varphi \mathbb{Z} \longrightarrow \mathfrak{A}$ such that ψ restricts to ρ on $C(X)$ and $\psi(U_\varphi) = V$. A construction of this C*-algebra can be found in [18] [12]. Given two compact separable spaces X and Y and two homeomorphisms φ and ψ of X and Y respectively, if φ and ψ are topologically conjugated in the sense that there exists an homeomorphism γ from X onto Y such that $\gamma \circ \varphi \circ \gamma^{-1} = \psi$, then by the universal property we have $C(X) \rtimes_\varphi \mathbb{Z} = C(Y) \rtimes_\psi \mathbb{Z}$. Thus, C*-crossed-products are invariants of topological conjugacy, though they are known not to be complete invariants.

1991 *Mathematics Subject Classification.* 46L05 (Primary), 37E30, 30D05 (Secondary).

Key words and phrases. conformal automorphisms of the disk, crossed products C*-algebras, Irreducible representations.

We study in this paper the case when $X = \mathbb{D} = \{z \in \mathbb{C} : |z| \leq 1\}$ and φ is a conformal automorphism of $\mathbb{D}$. The C^* -algebra $C(\mathbb{D})$ can be represented as the C^* -algebra $C^*(A_\varphi)$ for any normal element A_φ on some Hilbert space with $\sigma(A_\varphi) = \mathbb{D}$ where $\sigma(\cdot)$ denotes the spectrum of an operator. We denote the canonical unitary of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ by U_φ . Thus, $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ is the universal C^* -algebra generated by a unitary U_φ and a normal element A_φ such that $\sigma(A_\varphi) = \mathbb{D}$ and $U_\varphi^* A_\varphi U_\varphi = \varphi(A_\varphi)$. Any pair (A, U) where A is a normal operator with $\sigma(A) \subseteq \mathbb{D}$ and U a unitary operator on some Hilbert space with $U^* A U = \varphi(A)$ will be called a covariant pair for $(C(\mathbb{D}), \varphi, \mathbb{Z})$. Since $\sigma(A)$ is a compact subset of $\sigma(A_\varphi) = \mathbb{D}$, we define a $*$ -epimorphism ρ from $C(\mathbb{D})$ onto $C(\sigma(A))$ by setting $\rho(f(A_\varphi)) = f(A)$ for any $f \in C(\mathbb{D})$. We observe that $U^* f(A) U = f \circ \varphi(A) = f(U_\varphi^* A_\varphi U_\varphi)$, so by universality there exists a unique $*$ -epimorphism $\psi : C(\mathbb{D}) \rtimes_\varphi \mathbb{Z} \twoheadrightarrow C^*(A, U)$ such that $\psi(A_\varphi) = A$ and $\psi(U_\varphi) = U$.

While C^* -crossed products of minimal dynamical systems have received a lot of attention, more general and natural situations have not been studied as much. For crossed-product for action on the circle, one can consult [5]. We offer in this paper to investigate a few concrete examples of non-proper, non-free, non-minimal dynamical systems. Some of our examples are of type I, and some general properties of such crossed-products were derived in [16].

We start this paper with a complete classification of the conformal automorphisms of $\mathbb{D}$ up to topological conjugacy. We split the group $\text{M}_\mathbb{D}$ of conformal automorphisms of $\mathbb{D}$ into the set of hyperbolic automorphisms, which are all topologically conjugated with one another, the set of parabolic automorphisms which are all conformally conjugated with each other, the set of elliptic automorphisms, each of which is conjugated to a nontrivial rotation around the origin, and the identity map, which is a special case whose crossed-product is trivially $C(\mathbb{D}) \rtimes_{\text{Id}} \mathbb{Z} = C(\mathbb{D} \times \mathbb{T})$.

We then study the full C^* -crossed-product algebra for each conformal automorphism. In each case, we provide a full description of its Gelfand's spectrum, with its Zarisky (or hull-kernel) topology. We then describe all the irreducible representations, explicitly for the hyperbolic and parabolic cases, or based upon the representation theory of the rotation algebras for the elliptic case. Last, we derive an explicit description of each crossed-product in term of more elementary algebras, exposing the topological content of the C^* -crossed-product in each case. We prove that:

- For hyperbolic automorphisms, the crossed-product is isomorphic to

$$\left\{ \begin{array}{l} f \in C\left(\mathbb{R} \times [0, 1], \left[\begin{array}{cc} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{array} \right] \right) \text{ such that:} \\ \forall x \in \mathbb{R} \ \forall y \in [0, 1] \quad \left\{ \begin{array}{l} f(x, y) - f(0, 0) \in \mathcal{K} \\ U^* f(x, y) U = f(x + 1, y) \end{array} \right. \end{array} \right\},$$

where $\mathcal{T}$ is the Toeplitz algebra and $\mathcal{K}$ is the compact operator algebra,

- For parabolic automorphisms, the crossed-product is isomorphic to $C(U') + C_1(\mathbb{D}, \mathcal{K})$ where $U' : t \in \mathbb{D} \mapsto U$ with U the bilateral shift on $l^2(\mathbb{Z})$ and $\mathcal{K}$ the compact operator algebras acting on $l^2(\mathbb{Z})$,
- For an elliptic automorphism conjugated to the rotation of angle $2\pi\theta$, the crossed-product is isomorphic to

$$\{f \in C([0, 1], \mathcal{A}_\theta) : f(0) \in C^*(U)\},$$

where $\mathcal{A}_\theta = C^*(U, V)$ is the universal C^* -algebra generated by two unitaries U, V such that $VU = e^{2i\pi\theta}UV$.

In this paper, unless otherwise stated, the norm of a Banach space E is denoted by $\|\cdot\|_E$.

2. CLASSIFICATION OF THE CONFORMAL AUTOMORPHISMS OF THE CLOSED UNIT DISK

The following well-known corollary of the Schwarz lemma can be found in many textbooks in complex analysis (e.g. [9, Theorem 2.1 p.213]):

Proposition 2.1. *Let φ be a conformal automorphism of the closed unit disk $\mathbb{D} = \{z \in \mathbb{C} : |z| \leq 1\}$. Then there exists $z_0 \in \mathbb{C}$ and $\theta \in [0, 1)$ such that $|z_0| < 1$ and:*

$$\varphi(z) = \exp(2i\pi\theta) \frac{z - z_0}{1 - \overline{z_0}z}$$

for all $z \in \mathbb{D}$.

Moreover, $\varphi(0) = 0$ if and only if there exists $\theta \in [0, 1)$ such that for all $z \in \mathbb{D}$ we have $\varphi(z) = \exp(2i\pi\theta)z$.

We recall the following well-known definitions:

Definition 2.2. *Let ψ be an homeomorphism of $\mathbb{D}$. A point $\omega \in \mathbb{D}$ is attractive for the dynamical system $(C(\mathbb{D}), \psi, \mathbb{Z})$ when there exists an open neighborhood Ω of ω in $\mathbb{D}$ such that for all $y \in \Omega$ we have:*

$\lim_{n \rightarrow \infty} \psi^n(y) = \omega$. A point α is repulsive when there exists an open neighborhood Ω of α in $\mathbb{D}$ such that $\lim_{n \rightarrow \infty} \psi^{-n}(y) = \alpha$ for all $y \in \Omega$.

We now classify all the conformal automorphisms of $\mathbb{D}$ up to topological conjugacy, since C^* -crossed-products only depend on topological conjugacy classes. As a first step, we identify the conformal conjugacy classes of conformal automorphisms of $\mathbb{D}$.

Theorem 2.3. *Let $\varphi \in \mathbb{M}_{\mathbb{D}}$ where $\mathbb{M}_{\mathbb{D}}$ is the group of conformal automorphisms of $\mathbb{D}$: thus $\varphi : z \in \mathbb{D} \mapsto \exp(2i\pi\theta) \frac{z-z_0}{1-\bar{z}_0z}$ for some $\theta \in [0, 1)$ and $z_0 \in \mathbb{C}$ such that $|z_0| < 1$. Then:*

(1) *The following are equivalent:*

- $|z_0| > |\sin(\pi\theta)|$,
- φ has exactly two distinct fixed points in $\mathbb{D}$,
- φ has exactly two fixed points on $\mathbb{T}$,
- φ is conjugated in $\mathbb{M}_{\mathbb{D}}$ with $z \in \mathbb{D} \mapsto \frac{z+a}{1+\bar{a}z}$ for some unique $a \in (0, 1)$.

If these assertions hold, then φ is called hyperbolic and one fixed point of φ is attractive, while the other one is repulsive. In particular, $z \in \mathbb{D} \mapsto \frac{z+a}{1+\bar{a}z}$ for $a \in (0, 1)$ admits 1 as attractive fixed point and -1 as repulsive fixed point.

(2) *The following are equivalent:*

- $|\sin(\pi\theta)| > |z_0|$,
- φ has exactly one fixed point α in $\mathbb{D}$ such that $|\alpha| < 1$,
- φ is conjugated in $\mathbb{M}_{\mathbb{D}}$ to a rotation $z \in \mathbb{D} \mapsto \mu z$ for some unique $\mu \in \mathbb{T} \setminus \{1\}$.

If these assertions hold, then φ is called elliptic.

(3) *The following are equivalent:*

- $|\sin(\pi\theta)| = |z_0| \neq 0$,
- φ has exactly one fixed point α in $\mathbb{D}$ such that $|\alpha| = 1$,
- if $\varphi(-1)$ is on the upper half of $\mathbb{T}$, then φ is conjugated in $\mathbb{M}_{\mathbb{D}}$ with $\phi : z \mapsto \frac{3i-1}{i-3} \frac{z-\frac{2i-1}{i-3}}{1-\frac{-2i-1}{-i-3}z}$, where ϕ is the unique conformal automorphism of $\mathbb{D}$ whose only fixed point is 1 and $\phi(-1) = i$.
- If $\varphi(-1)$ is on the lower half of $\mathbb{T}$, then φ is conjugated in $\mathbb{M}_{\mathbb{D}}$ with $\phi : z \mapsto \frac{3i+1}{i+3} \frac{z-\frac{2i+1}{i+3}}{1+\frac{2i-1}{-i+3}z}$, where ϕ is the unique conformal automorphism of $\mathbb{D}$ whose only fixed point is 1 and $\phi(-1) = -i$.

If these assertions hold, φ is called parabolic and its fixed point is both attractive and repulsive.

- (4) Last, φ is the identity map if and only if it has more than two fixed points, if and only if $\theta = 0$ and $z_0 = 0$, if and only if it has two fixed points α, β with either $|\alpha| = 1, |\beta| < 1$ or $|\alpha| < 1, |\beta| < 1$.

Proof. The fixed points of φ are the solutions of $\overline{z_0}z^2 + (e^{2i\pi\theta} - 1)z - e^{2i\pi\theta}z_0 = 0$. Thus, by a straightforward computation, the location of the fixed points of $\varphi \neq \text{Id}$ are determined by the sign of $|z_0| - |\sin(\pi\theta)|$ as stated in the theorem (see appendix at the end for the calculation).

We now turn to the matter of proving that φ is, in each case, conjugated to a specific conformal automorphism. First, let us assume that φ has two distinct fixed points $\alpha, \beta \in \mathbb{T}$. Assume first $\alpha, \beta \notin \{-1, 1\}$. Let ω be a square root of $\overline{\alpha\beta}$ and let $\gamma = \frac{i-\omega\alpha}{1-i\omega\alpha}$ (observe that $\gamma \in]-1, 1[$). We set $\psi(z) = i\omega \frac{z+\gamma\overline{\omega}}{1+\gamma\omega z}$ for all $z \in \mathbb{D}$. Then $\psi \circ \varphi \circ \psi^{-1}$ fixes -1 and 1 .

Now, any conformal automorphism φ has -1 and 1 as fixed points if and only if $\lambda - 1 = 0$ and $z_0/\overline{z_0} = 1$, namely $z_0 \in \mathbb{R}$ and $e^{2i\pi\theta} = 1$. Up to conjugating φ by $\psi' : z \mapsto -z$ we can assume $z_0 \leq 0$, so all hyperbolic automorphisms of $\mathbb{D}$ are conjugated in $\mathbb{M}_{\mathbb{D}}$ to $z \mapsto \frac{z+z_0}{1+z_0z}$ for some $z_0 \in (0, 1)$, which a simple calculation shows to be uniquely determined by φ (see appendix at the end for the calculation).

If instead $\{\alpha, \beta\} \not\subset \mathbb{T}$ then φ has only one fixed point $c \in \{\alpha, \beta\}$ in $\mathbb{D}$. Then set $\psi : z \mapsto \frac{z-c}{1-\overline{c}z} \in \mathbb{M}_{\mathbb{D}}$. The map $R = \psi \circ \varphi \circ \psi^{-1}$ is a conformal automorphism of $\mathbb{D}$ with $R(0) = 0$. Hence R is a rotation by Theorem (2.1). Now, if $\varphi \in \mathbb{M}_{\mathbb{D}}$ conjugates $z \mapsto \mu z$ with $z \mapsto \nu z$ for $\mu, \nu \in \mathbb{T}$ then $\varphi(0) = 0$ so $\varphi : z \mapsto \eta z$ for $\eta \in \mathbb{T}$ and thus $\mu = \nu$.

Last, observe that when φ is parabolic with fixed point α then $r_{\alpha^{-1}} \circ \varphi \circ r_{\alpha}(1) = 1$, where $r_{\alpha} : z \mapsto \alpha z$. Hence up to conformal conjugacy, we may as well assume that φ admits 1 as its unique fixed point. Let z_0 with $|z_0| < 1$ and $\lambda \in \mathbb{T}$ such that $\varphi : z \mapsto \lambda \frac{z-z_0}{1-\overline{z_0}z}$. Now

$\varphi(1) = 1$ if and only if $1 = \frac{1-\lambda}{2\overline{z_0}}$, so if and only if $\varphi : z \mapsto \lambda \frac{z-\frac{1-\lambda}{2}}{1-\frac{1-\lambda}{2}\overline{z_0}z}$. Now,

if $\varphi : z \mapsto \lambda \frac{z-\frac{1-\lambda}{2}}{1-\frac{1-\lambda}{2}\overline{z_0}z}$ for some $\lambda \in \mathbb{T}$, then let ψ be the (unique) conformal automorphism of $\mathbb{D}$ such that $\psi(1) = 1, \psi(-1) = -1$ and $\psi(\varphi(-i)) = i$ if $\varphi(-i)$ is in the upper half plane, or $\psi(\varphi(-1)) = -i$ otherwise. Then $\phi = \psi \circ \varphi \circ \psi^{-1}$ is parabolic since it has a unique fixed point in $\mathbb{T}$ and we have $\phi(1) = 1$ and $\phi(-1) = i$ if $\varphi(-1)$ is in the upper half plane, and $\phi(-1) = -i$ otherwise. We deduce that $\phi : z \mapsto \frac{3i-1}{i-3} \frac{z-\frac{2i-1}{i-3}}{1-\frac{2i-1}{-i-3}\overline{z_0}z}$ (if $\varphi(-1)$

is in the upper half-plane) or $\phi : z \mapsto \frac{3i+1}{i+3} \frac{z-\frac{2i+1}{i+3}}{1+\frac{2i-1}{-i+3}\overline{z_0}z}$. This concludes our proof. $\square$

Now, we have reduced the topological conjugacy classification problem to classifying the special automorphisms in Proposition (2.3):

Theorem 2.4. *Let φ, ψ be two conformal automorphisms of $\mathbb{D}$. Then:*

- *If φ and ψ are topologically conjugated and if φ is hyperbolic (resp. parabolic, elliptic) then ψ is hyperbolic (resp. parabolic, elliptic),*
- *If φ is elliptic then φ is topologically conjugated to a rotation $r_\theta : z \in \mathbb{D} \mapsto e^{2i\pi\theta}z$ for some $\theta \in (0, 1)$. Moreover, r_θ and $r_{\theta'}$ are topologically conjugated if and only if $\theta \in \{\theta', -\theta'\}$.*
- *All parabolic automorphisms are topologically conjugated on $\mathbb{D}$,*
- *All hyperbolic automorphisms are topologically conjugated on $\mathbb{D}$. Moreover, if $\varphi : z \mapsto \frac{z+a}{1+az}$ for some $a \in (0, 1)$ and if we set:*

$$\mathcal{D}_\varphi = \{z \in \mathbb{D} : \operatorname{Re}(z) \geq 0, |1 - z| \geq 1 - a\}$$

then for any $x \in \mathbb{D} \setminus \{-1, 1\}$ there exists a unique $y \in \mathcal{D}_\varphi$ such that $\mathcal{O}_\varphi(x) = \mathcal{O}_\varphi(y)$, where $\mathcal{O}_\varphi(w) = \{\varphi^n(w) : n \in \mathbb{Z}\}$ is the orbit for φ of any $w \in \mathbb{D}$.

Proof. It is well known that two rotations $r_\lambda : z \mapsto \lambda z$ and $r_\mu : z \mapsto \mu z$ for $\lambda, \mu \in \mathbb{T}$ are conjugated if and only if $\lambda \in \{\mu, \bar{\mu}\}$: if $\lambda = \mu$ it is trivial, if $\lambda = \bar{\mu}$ then $\overline{(r_\lambda(\bar{z}))} = r_\mu$, so the condition is sufficient, and it is necessary by [8, Theorem 11.2.7, p. 397]. Since the number of fixed points is a topological invariant, and any homeomorphism of $\mathbb{D}$ maps $\mathbb{T}$ to $\mathbb{T}$, the two first statements of our theorem are proven.

By Theorem (2.3), if φ is parabolic then it is conformally, hence topologically conjugated to either

$$\phi_+ : z \mapsto \frac{3i-1}{i-3} \frac{z - \frac{2i-1}{i-3}}{1 - \frac{-2i-1}{-i-3}z} \quad \text{or} \quad \phi_- : z \mapsto \frac{3i+1}{i+3} \frac{z - \frac{2i+1}{i+3}}{1 + \frac{2i-1}{-i+3}z}.$$

Yet, if $\nu : z \mapsto \bar{z}$ then $\nu^{-1} \circ \phi_- \circ \nu = \phi_+$, hence the third assertion of our theorem.

By Proposition (2.3), any hyperbolic automorphism f is conformally, hence topologically conjugated in $\mathbb{D}$ to an hyperbolic automorphism φ with fixed points -1 and 1 where 1 is the attractive fixed point. It suffices therefore to prove that any two hyperbolic automorphisms φ and ψ of $\mathbb{D}$ with fixed points -1 and 1 , where 1 is attractive, are topologically conjugated. Therefore, let $\varphi : z \mapsto \frac{z+a}{1+az}$ and $\psi : z \mapsto \frac{z+b}{1+bz}$ for $a, b \in (0, 1)$.

We define the region $\overline{\mathcal{D}_\varphi}$ as the closed subset of $\mathbb{D}$ contained between the diameter $L_\varphi = \{it : t \in [-1, 1]\}$ and its image $\varphi(L)$, which is the circle of center $\varphi(\infty) = \frac{1}{a}$ and radius $\frac{1}{a} - a$. We denote by $\mathcal{D}_\varphi$ the

set $\overline{\mathcal{D}_\varphi} \setminus \varphi(L)$ and we check easily that the orbit for φ of any point of $\mathbb{D} \setminus \{-1, 1\}$ intersects $\mathcal{D}_\varphi$ at exactly one point.

The region $\overline{\mathcal{D}_\varphi}$ is homeomorphic to the rectangle $[0, 1]^2$. We choose an homeomorphism $\mu_\varphi : \overline{\mathcal{D}_\varphi} \rightarrow [0, 1]^2$ such that $\mu_\varphi(it) = (t, 0)$ and $\mu_\varphi(\varphi(it)) = (t, 1)$ for all $t \in [-1, 1]$.

Of course, we can as well construct an homeomorphism $\mu_\psi : \overline{\mathcal{D}_\psi} \rightarrow [0, 1]^2$ such that $\mu_\psi(it) = (t, 0)$ and $\mu_\psi(\psi(it)) = (t, 1)$. Hence $\mu = \mu_\psi^{-1} \mu_\varphi$ is an homeomorphism from $\overline{\mathcal{D}_\varphi}$ onto $\overline{\mathcal{D}_\psi}$ such that $\mu \circ \varphi = \psi \circ \mu$ on the set $\{it : t \in [-1, 1]\}$. By a trivial induction, we can extend μ to the set $\mathbb{D} \setminus \{-1, 1\}$ and then set $\mu(-1) = -1$ and $\mu(1) = 1$, and this map is easily checked to be an homeomorphism of $\mathbb{D}$ such that $\mu \circ \varphi = \psi \circ \mu$ on $\mathbb{D}$ (see appendix for a more explicit construction of an example of μ). $\square$

This achieves the classification, up to topological classification, of all the conformal automorphisms of $\mathbb{D}$. In particular, the description of the dynamical system $(C(\mathbb{D}), \varphi, \mathbb{Z})$ is now easy. If φ is hyperbolic, then (up to conformal conjugacy) one has -1 as a repulsive fixed point, 1 as attractive fixed point, and one can easily check that every orbit besides $\{1\}$ and $\{-1\}$ for φ is infinite, discrete and supported on a circle passing by the points -1 and 1 . If φ is elliptic, then up to topological equivalence φ is simply a rotation around the origin. If φ is parabolic, it has a unique fixed point 1 (up to conformal conjugacy) which is both attractive and repulsive, and its orbits besides $\{1\}$ are infinite, discrete and supported on circles tangent at 1 to the vertical line of equation $x = 1$.

3. HYPERBOLIC AUTOMORPHISMS

Let φ be a hyperbolic automorphism of $\mathbb{D}$ which fixes 1 and -1 . By Theorem (2.4), any hyperbolic automorphism ψ of $\mathbb{D}$ is topologically conjugated to φ . Therefore, the C^* -crossed-products $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ and $C(\mathbb{D}) \rtimes_\psi \mathbb{Z}$ are $*$ -isomorphic. This section describes the crossed-product $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$.

The orbit space of φ restricted to $\mathbb{D} \setminus \{-1, 1\}$ is denoted $\mathcal{C}$ and, topologically, is the cylinder $[0, 1] \times \mathbb{T}$. There is a one-to-one continuous map $\mathcal{D}_\varphi \rightarrow \mathcal{C}$, whose inverse is not continuous (note that $\mathcal{C}$ is compact while $\mathcal{D}_\varphi$ is not).

3.1. Irreducible representations. We start with a full description of the irreducible $*$ -representations of the dynamical system $(C(\mathbb{D}), \varphi, \mathbb{Z})$. We first establish a result on the representations of general discrete

dynamics $(C(\mathbb{D}), \psi, \mathbb{Z})$. We will denote the orbit for ψ of any $x \in \mathbb{D}$ by $\mathcal{O}_\psi(x) = \{\psi^z(x) : z \in \mathbb{Z}\}$.

Lemma 3.1. *Let ψ be any homeomorphism of the closed unit disk $\mathbb{D}$. Let π be an irreducible representation of $C(\mathbb{D}) \rtimes_\psi \mathbb{Z}$ on some Hilbert space $\mathcal{H}$. Then $\sigma(\pi(A_\psi))$ is a ψ -invariant compact subset of $\mathbb{D}$. Moreover, for any ψ -invariant Borel subset E of $\sigma(\pi(A_\psi))$, the spectral projection of $\pi(A_\psi)$ associated to E is either 0 or 1. Hence, if E is a nonempty relatively open invariant subset of $\sigma(A_\psi)$ then $\sigma(A_\psi) = \overline{E}$.*

Proof. Let $A_\pi = \pi(A_\psi)$ and $U_\pi = \pi(U_\psi)$. We have $U_\pi^* A_\pi U_\pi = \psi(A_\pi)$, so by the spectral mapping theorem [3, Theorem 8.11, p.289], $\sigma(A_\pi)$ is a ψ -invariant set. Moreover, if we denote by χ_E the characteristic function of any Borel subset E of $\sigma(A_\pi)$, then by the spectral theorem we have:

$$U_\pi^* \chi_E(A_\pi) U_\pi = \chi_E(U_\pi^* A_\pi U_\pi) = \chi_{\psi^{-1}(E)}(A_\pi).$$

In particular, assume E is ψ -invariant. Then the spectral projection $\chi_E(A_\pi)$ of A_π commutes with U_π (and with A_π by definition), hence since π is irreducible we deduce that $\chi_E \in \{0, 1\}$. If E is relatively open in $\sigma(A_\pi)$ and nonempty, then $\chi_E(A_\pi) > 0$, so for any nonempty relatively open ψ -invariant subset of $\sigma(A_\pi)$ we have $\chi_E(A_\pi) = 1$. In this case, $\chi_{\overline{E}}(A_\pi) = 1$, so $\chi_{\sigma(A_\pi) \setminus \overline{E}}(A_\pi) = 0$. Yet $\sigma(A_\pi) \setminus \overline{E}$ is relatively open in $\sigma(A_\pi)$, so by the spectral theorem $\sigma(A_\pi) \setminus \overline{E} = \emptyset$. $\square$

Theorem 3.2. *Let ψ be an homeomorphism of the closed unit disk $\mathbb{D}$. Let π be an irreducible $*$ -representation of $C(\mathbb{D}) \rtimes_\psi \mathbb{Z}$. Then there exists $x \in \mathbb{D}$ such that the spectrum $\sigma(\pi(A_\psi))$ of $\pi(A_\psi)$ is $\overline{\mathcal{O}_\psi(x)}$.*

Proof. Let $A_\pi = \pi(A_\psi)$. For any subset E of $\sigma(A_\pi)$, we denote the set $\bigcup_{n \in \mathbb{Z}} \psi^n(E)$ by $\mathcal{O}_\psi(E)$.

Let $(\Omega_n)_{n \in \mathbb{N}}$ be a countable basis of relatively open nonempty subsets of $\sigma(A_\pi)$ for the topology of the second countable compact space $\sigma(A_\pi)$. Let Θ be a relatively open, nonempty subset of $\sigma(A_\pi)$ and let $n \in \mathbb{N}$. By Lemma (3.1), we have $\sigma(A_\pi) = \overline{\mathcal{O}_\psi(\Omega_n)}$, so $\Theta \cap \mathcal{O}_\psi(\Omega_n) \neq \emptyset$. Now since $\sigma(A_\pi)$ is a compact Hausdorff space, there exists a nonempty open set V such that $\overline{V} \subseteq \Theta$, and we can choose $V \subseteq \mathcal{O}_\psi(\Omega_n)$ for any choice of $n \in \mathbb{N}$ by Lemma (3.1). We can therefore construct by induction a sequence $(V_n)_{n \in \mathbb{N}}$ of relatively open nonempty subsets of $\sigma(A_\pi)$ with $V_0 = \Omega_0$ and such that for all $n \in \mathbb{N}$ we have $\overline{V_{n+1}} \subseteq V_n$ and $V_n \subseteq \mathcal{O}_\psi(\Omega_n)$. Therefore, as $\sigma(A_\pi)$ is compact, $\bigcap_{n \in \mathbb{N}} V_n = \bigcap_{n \in \mathbb{N}} \overline{V_n}$ is

nonempty and by construction $\bigcap_{n \in \mathbb{N}} V_n \subseteq \bigcap_{n \in \mathbb{N}} \mathcal{O}_\psi(\Omega_n)$. Let $x \in \bigcap_{n \in \mathbb{N}} V_n$: then since $x \in \bigcap_{n \in \mathbb{N}} \mathcal{O}_\psi(\Omega_n)$, we have $\mathcal{O}_\psi(x) \cap \mathcal{O}_\psi(\Omega_n) \neq \emptyset$ for all $n \in \mathbb{N}$. Therefore $\mathcal{O}_\psi(x) \cap \Omega_n \neq \emptyset$. Thus, we have $\mathcal{O}_\psi(x) \cap \Theta \neq \emptyset$ since $(\Omega_n)_{n \in \mathbb{N}}$ is a basis for the topology of $\sigma(A_\pi)$. Hence $\overline{\mathcal{O}_\psi(x)} = \sigma(A_\pi)$. $\square$

A way to read Lemma (3.1) is to say that $\sigma(\pi(A_\varphi))$ is topologically transitive for any irreducible representation π . Theorem (3.2) holds in greater generality, for any second countable Hausdorff compact space instead of $\mathbb{D}$, with essentially the same proof. We only consider the case of the disk for our paper.

We now work with our hyperbolic automorphism φ of $\mathbb{D}$. To fully describe the image of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ by its irreducible representations, we will find the following lemmas useful. We start by introducing some terminology: we let $\overline{\mathbb{Z}} = \mathbb{Z} \cup \{-\infty, \infty\}$, and we let $\mathbb{Z}$ acts by translation on $\overline{\mathbb{Z}}$. More precisely, let τ be the action of $\mathbb{Z}$ on $\overline{\mathbb{Z}}$ defined by $\tau_z(z') = z + z'$, where $z + \infty = \infty$ and $z - \infty = -\infty$ for all $z \in \mathbb{Z}$ and $z' \in \overline{\mathbb{Z}}$. We define:

Definition 3.3. Let $\mathbb{H} = l^2(\mathbb{Z})$ and let $(e_n)_{n \in \mathbb{Z}}$ be the canonical Hilbert basis of $\mathbb{H}$. We define:

$$C_{0,0}(\mathbb{Z}) = \left\{ (\lambda_n)_{n \in \mathbb{Z}} : \lim_{n \rightarrow \infty} \lambda_n = \lim_{n \rightarrow -\infty} \lambda_n = 0 \right\},$$

$$C(\overline{\mathbb{Z}}) = \left\{ (\lambda_n)_{n \in \mathbb{Z}} : (\lambda_n)_{n \in \mathbb{N}} \text{ and } (\lambda_{-n})_{n \in \mathbb{N}} \text{ converge} \right\}.$$

For any $s = (\lambda_n)_{n \in \mathbb{Z}} \in C(\overline{\mathbb{Z}})$, let $\text{Diag}(s)$ be the unique bounded linear operator on $\mathbb{H}$ such that $\text{Diag}(s)(e_n) = \lambda_n e_n$ for all $n \in \mathbb{Z}$. We set $\mathfrak{D} = \text{Diag}(C(\overline{\mathbb{Z}}))$ and $\mathfrak{D}_{0,0} = \text{Diag}(C_{0,0}(\mathbb{Z}))$. Of course, $\mathfrak{D}$ and $\mathfrak{D}_{0,0}$ are C^* -algebras, and Diag is a $*$ -isomorphism between $C(\overline{\mathbb{Z}})$ and $\mathfrak{D}$ and between $\mathfrak{D}_{0,0}$ and $C_{0,0}(\mathbb{Z})$.

Definition 3.4. We define U on $\mathbb{H} = l^2(\mathbb{Z})$ by $U(\xi_n)_{n \in \mathbb{N}} = (\xi_{n-1})_{n \in \mathbb{N}}$ for all $\xi \in \mathbb{H}$. Namely, U is the bilateral shift unitary operator on $\mathbb{H}$.

Definition 3.5. We endow $\overline{\mathbb{Z}}$ with the topology of the Gelfand's spectrum of $C(\overline{\mathbb{Z}})$ (this is of course the usual topology on $\overline{\mathbb{Z}}$).

The next two lemmas describe the crossed-product $C(\overline{\mathbb{Z}}) \rtimes_\tau \mathbb{Z}$, which can be seen as the crossed-product $C(\overline{\mathcal{O}_\varphi(x)}) \rtimes_\varphi \mathbb{Z}$ when $x \in \mathbb{D} \setminus \{-1, 1\}$ and is therefore a fundamental building block of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$. Let $\mathcal{T}$ be the Toeplitz C^* -algebra $C^*(S)$ where S is the unilateral shift on $l^2(\mathbb{N})$ defined by $S(x_0, x_1, \dots) = (0, x_0, x_1, \dots)$ for all $(x_n)_{n \in \mathbb{N}} \in l^2(\mathbb{N})$.

Lemma 3.6. *Let $E = \{U^n \mathfrak{D}_{0,0} : n \in \mathbb{Z}\}$. The C^* -algebra $C^*(E)$ is the C^* -algebra of compact operators $\mathcal{K}$ and the C^* -algebra $C^*(U, \mathfrak{D})$ equals $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$ where $\mathbb{H} = \mathbb{H}_- \oplus \mathbb{H}_+$ with $\mathbb{H}_- = \{e_n : n < 0\}$, $\mathbb{H}_+ = \{e_n : n \geq 0\}$ and with the natural identification $\mathbb{H}_+ = \mathbb{H}_- = l^2(\mathbb{N})$. Moreover, if χ is defined by the exact sequence $0 \rightarrow \mathcal{K} \rightarrow \mathcal{T} \xrightarrow{\chi} C(\mathbb{T}) \rightarrow 0$ and $\gamma = \chi \oplus \chi$ act on $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$, then $\gamma(\text{Diag}(f)U^n) = f(-\infty)Z^{-n} \oplus f(\infty)Z^n$ for all $f \in C(\overline{\mathbb{Z}})$, $n \in \mathbb{Z}$ and where $Z : z \in \mathbb{T} \mapsto z$.*

Thus, $A \in \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$ if and only if there exists $f, g \in C(\mathbb{T})$ and a compact operator k on $\mathbb{H}$ such that for any $\xi_+ \oplus \xi_- \in \mathbb{H}_- \oplus \mathbb{H}_+$ we have $A\xi = (T_f\xi_+ + T_g\xi_-) + k(\xi_+ + \xi_-)$ where T_f and T_g are the Toeplitz operators of symbols f and g .

Proof. By construction, $\mathfrak{D}_{0,0} \subset \mathcal{K}$, so $C^*(E) \subseteq \mathcal{K}$ as $\mathcal{K}$ is an ideal in $\mathcal{B}(\mathbb{H})$. Let now P be any projection in $\mathcal{B}(\mathbb{H})$. Assume that P commutes with $C^*(E)$. Then P commutes with $\mathfrak{D}_{0,0}$ so P is diagonal in the basis $(e_n)_{n \in \mathbb{Z}}$. If $P \notin \{0, 1\}$ then there exists $k \in \mathbb{Z}$ such that $Pe_k = 0$ but $Pe_{k-1} = e_{k-1}$. Let Q be the projection on $\text{span}\{e_n : |n| \leq k\} \in \mathfrak{D}_{0,0}$. Now, by assumption $UQP = PUQ$ yet $UQPe_{k-1} = e_k$ but $PUe_{k-1} = 0$. This is a contradiction, so $P \in \{0, 1\}$. Hence $C^*(E) = \mathcal{K}$ since it is irreducible [1, Theorem 1.4.2, p. 18].

Write $U = \begin{bmatrix} U_{11} & 0 \\ U_{21} & U_{22} \end{bmatrix}$ according to the decomposition $l^2(\mathbb{Z}) = \mathbb{H}_- \oplus \mathbb{H}_+$. Now, $U_{21} \in \mathcal{K}$ as it is the rank one operator $U_{21} = \langle \cdot, e_{-1} \rangle e_0$ (where $\langle \cdot, \cdot \rangle$ is the inner product of $\mathbb{H}$). With the obvious identifications of $\mathbb{H}_-$ and $\mathbb{H}_+$ with $l^2(\mathbb{N})$ it is immediate that $U_{11} = S^*$ and $U_{22} = S$. Now, let A be the projection on $\mathbb{H}_+$: by construction, $A \in \mathfrak{D}$ and $AUA = U_{22} \in C^*(U, \mathfrak{D})$. Similarly, $U_{11} \in C^*(U, \mathfrak{D})$, so $\mathcal{T} \oplus \mathcal{T} \subseteq C^*(U, \mathfrak{D})$. Since $C^*(E) \subset C^*(U, \mathfrak{D})$ we deduce that $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix} \subseteq$

$C^*(U, \mathfrak{D})$ (observe that $\mathcal{K}$ is $\begin{bmatrix} \mathcal{K} & \mathcal{K} \\ \mathcal{K} & \mathcal{K} \end{bmatrix}$ in our decomposition of $l^2(\mathbb{Z})$).

On the other hand, by definition $C^*(U, \mathfrak{D})$ is the smallest C^* -algebra containing U and $\mathfrak{D}$. Since $U \in \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$ and $\mathfrak{D} \subset \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$ we

deduce that $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix} = C^*(U, \mathfrak{D})$. As a last remark, Let $f \in C(\overline{\mathbb{Z}})$.

Then $\text{Diag}(f) = (1 - A)\text{Diag}(f) \oplus A\text{Diag}(f)$ and $A\text{Diag}(f) = f(\infty)1 + A(\text{Diag}(f) - f(\infty)1)$ where now $A(\text{Diag}(f) - f(\infty)1) \in \mathcal{K}$. Similarly,

$(1 - A) \text{Diag}(f) = f(-\infty)1 + (1 - A)(\text{Diag}(f) - f(-\infty)1)$ and $(1 - A)(\text{Diag}(f) - f(-\infty)1) \in \mathcal{K}$. Since $\gamma(\mathcal{K}) = \{0\}$ and γ is a $*$ -morphism with $\gamma(U) = Z^* \oplus Z$, we deduce that $\gamma(\text{Diag}(f)U^n) = f(-\infty)Z^{-n} \oplus f(\infty)Z^n$. $\square$

Lemma 3.7. *The C^* -algebra $C(\overline{\mathbb{Z}}) \rtimes_{\tau} \mathbb{Z}$ is $*$ -isomorphic to $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$.*

Proof. Since $\mathbb{Z}$ is τ -invariant, we have by [10, Proposition 2.8]:

$$0 \longrightarrow C_{0,0}(\mathbb{Z}) \rtimes_{\tau} \mathbb{Z} \longrightarrow C(\overline{\mathbb{Z}}) \rtimes_{\tau} \mathbb{Z} \xrightarrow{\gamma'} C(\{-\infty, \infty\}) \rtimes_{\tau} \mathbb{Z} \longrightarrow 0.$$

Of course $C(\{-\infty, \infty\}) \rtimes_{\tau} \mathbb{Z} = C(\mathbb{T}) \oplus C(\mathbb{T})$ since both ∞ and $-\infty$ are fixed point for τ . Moreover, the C^* -algebra $C_{0,0}(\mathbb{Z}) \rtimes_{\tau} \mathbb{Z}$ is $*$ -isomorphic to $\mathcal{K}$ [7, 8.4.3, p. 141] and is the C^* -algebra $C^*(C_{0,0}(\mathbb{Z})V^n)_{n \in \mathbb{Z}}$ where V is the canonical unitary in $C(\overline{\mathbb{Z}}) \rtimes_{\tau} \mathbb{Z}$.

Let us now define a map from $C(\overline{\mathbb{Z}}) \rtimes_{\tau} \mathbb{Z}$ onto $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$ as follows. Let U be the bilateral shift in $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$. We set $\rho(V) = U$, and for $f \in C(\overline{\mathbb{Z}})$ we set $\rho(f) = \text{Diag}(f)$. By universality of the crossed-product $C(\overline{\mathbb{Z}}) \rtimes_{\tau} \mathbb{Z}$, we deduce that ρ extends into an epimorphism from $C(\overline{\mathbb{Z}}) \rtimes_{\tau} \mathbb{Z}$ onto $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$, since by construction $U^* \text{Diag}(f)U = \text{Diag}(g)$ where $g = f(\cdot + 1)$. Moreover, $\rho(fV^n) = \text{Diag}(f)U^n \in \mathcal{K}$ by construction, so $\rho(C_{0,0}(\mathbb{Z}) \rtimes_{\tau} \mathbb{Z}) = C^*(\{U^n \mathfrak{D}_{0,0} : n \in \mathbb{Z}\}) = \mathcal{K}$ by Lemma (3.6). Last, we observe that $\gamma'(\text{Diag}(f)V^n) = f(-\infty)Z^{\varepsilon_1 n} \oplus f(\infty)Z^{\varepsilon_2 n}$ for all $f \in C(\overline{\mathbb{Z}})$, $n \in \mathbb{Z}$, where $Z : z \in \mathbb{T} \mapsto z$ and with $\varepsilon_1, \varepsilon_2 \in \{-1, 1\}$. Up to a $*$ -automorphism of $C(\mathbb{T}) \oplus C(\mathbb{T})$ we may as well choose $\varepsilon_1 = -1$ and $\varepsilon_2 = 1$. On the other hand, $\gamma(\rho(\text{Diag}(f)V^n)) = f(-\infty)Z^{-n} \oplus f(\infty)Z^n$ by construction of ρ and Lemma (3.6). Hence, we have the following commuting diagram:

$$\begin{array}{ccccccc} 0 & \rightarrow & C_0(\mathbb{Z}) \rtimes_{\tau} \mathbb{Z} & \rightarrow & C(\overline{\mathbb{Z}}) \rtimes_{\tau} \mathbb{Z} & \rightarrow & C(\mathbb{T}) \oplus C(\mathbb{T}) \rightarrow 0 \\ & & \downarrow \rho & & \downarrow \rho & & \parallel \\ 0 & \rightarrow & \mathcal{K} & \rightarrow & \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix} & \rightarrow & C(\mathbb{T}) \oplus C(\mathbb{T}) \rightarrow 0 \end{array}$$

where the top and bottom lines are exact. Now, since $\mathcal{K}$ is simple, ρ in the first vertical arrow is injective, hence a $*$ -isomorphism. The last vertical arrow is just the identity of $C(\mathbb{T}) \oplus C(\mathbb{T})$. We deduce that ρ in the second vertical arrow is a $*$ -isomorphism by the Short Five Lemma [11, Lemma 3.1, p. 13]. $\square$

We now define two families of $*$ -representations of $C^*(U_\varphi, A_\varphi)$, which we shall then prove in Theorem (3.10) are the only ones up to unitary equivalence:

Definition 3.8. For any $\varepsilon \in \{-1, 1\}$ and any $\theta \in [0, 1)$ we define the representation $\pi_{\theta, \varepsilon}$ by $\pi_{\theta, \varepsilon}(A_\varphi) = \varepsilon$ and $\pi_{\theta, \varepsilon}(U_\varphi) = \exp(2i\pi\theta)$.

For any $x \in \mathbb{D} \setminus \{-1, 1\}$, we define the representation π_x on $l^2(\mathbb{Z})$ by letting $\pi_x(A_\varphi) = \text{Diag}((\varphi^n(x))_{n \in \mathbb{Z}})$ and $\pi_x(U_\varphi) = U$ for all $n \in \mathbb{Z}$.

We let $\text{IrrRep} = \{\pi_x, \pi_{\theta, \varepsilon} : x \in \mathbb{D} \setminus \{-1, 1\}, \varepsilon \in \{-1, 1\}, \theta \in [0, 1)\}$.

We indeed check that for any $x \in \mathbb{D} \setminus \{-1, 1\}$, $\theta \in [0, 1)$ and $\varepsilon \in \{-1, 1\}$, if $\pi = \pi_x$ or $\pi = \pi_{\theta, \varepsilon}$ then we have $\pi(U_\varphi^*)\pi(A_\varphi)\pi(U_\varphi) = \varphi(\pi(A_\varphi))$, so π extends to a $*$ -representation of $C^*(A_\varphi, U_\varphi)$ by universality. Now:

Lemma 3.9. All the representations in IrrRep are irreducible. Let $x, y \in \mathbb{D} \setminus \{-1, 1\}$, $\varepsilon, \varepsilon' \in \{-1, 1\}$ and $\theta, \theta' \in [0, 1)$.

We have $\pi_x(C^*(U_\varphi, A_\varphi)) = \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$, and moreover π_x is unitarily equivalent to π_y if and only if $\mathcal{O}_\varphi(x) = \mathcal{O}_\varphi(y)$, and $\pi_{\theta, \varepsilon}$ is unitarily equivalent to $\pi_{\varepsilon', \theta'}$ if and only if $(\varepsilon, \theta) = (\varepsilon', \theta')$.

Proof. For all representation π the spectra of $\pi(A_\varphi)$ and $\pi(U_\varphi)$ are invariant by unitary conjugation, it is also straightforward that no two distinct elements of IrrRep are unitarily conjugated, and that if π_x and π_y are conjugated for $x, y \in \mathbb{D} \setminus \{-1, 1\}$ then $\overline{\mathcal{O}_\varphi(x)} = \sigma(\pi_x(A_\varphi))$ equals to $\sigma(\pi_y(A_\varphi)) = \overline{\mathcal{O}_\varphi(y)}$, and so $\mathcal{O}_\varphi(x) = \overline{\mathcal{O}_\varphi(x)} \setminus \{-1, 1\} = \overline{\mathcal{O}_\varphi(y)} \setminus \{-1, 1\} = \mathcal{O}_\varphi(y)$. Conversely, if $\mathcal{O}_\varphi(x) = \mathcal{O}_\varphi(y)$ then there exists $n \in \mathbb{Z}$ such that $x = \varphi^n(y)$, so $U^{*n}\pi_x U^n = \pi_y$. The one-dimensional representations $\pi_{\theta, \varepsilon}$, where $\theta \in [0, 1)$ and $\varepsilon \in \{-1, 1\}$, are obviously irreducible. If $x \in \mathcal{D}_\varphi$ then, by Lemma (3.6), we have $\pi_x(C^*(U_\varphi, A_\varphi)) =$

$\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$, since $C^*(\pi_x(A_\varphi)) = \mathfrak{D}$ for all $x \in \mathbb{D} \setminus \{-1, 1\}$. As in Lemma (3.6), a projection P of $\mathbb{H}$ commutes with $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$ if and only if $P \in \{0, 1\}$ so π_x is irreducible. $\square$

Theorem 3.10. Let φ be an hyperbolic conformal automorphism of the closed unit disk $\mathbb{D}$. Let π be an irreducible representation for the dynamical system $(C(\mathbb{D}), \varphi, \mathbb{Z})$. Then π is unitarily equivalent to one of the representations in IrrRep .

Proof. Let π be an irreducible representation of $(C(\mathbb{D}), \varphi, \mathbb{Z})$ on some Hilbert space $\mathcal{H}$ with inner product $\langle \cdot, \cdot \rangle$. Let $A_\pi = \pi(A_\varphi)$ and $U_\pi =$

$\pi(U_\varphi)$. Then $\sigma(A_\pi) = \overline{\mathcal{O}_\varphi(x)}$ for some $x \in \mathbb{D}$ by Theorem (3.2). Now for our choice of automorphism φ , we observe that $x \in \overline{\mathcal{O}_\varphi(x)}$ is always isolated in $\overline{\mathcal{O}_\varphi(x)}$. Hence x is an eigenvalue for A_π . Let $\xi_x \in \mathcal{H}$ be a normalized eigenvector for A_π associated to x . Let $\mathcal{M} = \overline{\text{span}} \{U_\pi^n \xi_x : n \in \mathbb{Z}\}$. By construction, $\mathcal{M}$ is a reducing subspace for both U_π and A_π . Since π is irreducible, $\mathcal{M} \in \{\mathcal{H}, \{0\}\}$, yet since $\xi_x \neq 0$ we have $\mathcal{M} = \mathcal{H}$. Of course, $A_\pi U_\pi^n \xi_x = U_\pi^n \varphi^n(A_\pi) \xi_x = \varphi^n(x) U_\pi^n \xi_x$; in other words $U_\pi^n \xi_x$ is a normalized eigenvector for A_π associated to the eigenvalue $\varphi^n(x)$.

If $x \in \{-1, 1\}$ then $\varphi^n(x) = x$ for all $n \in \mathbb{Z}$. Hence $A_\pi = xI$ where I is the identity of $\mathcal{H}$. Therefore, $C^*(A_\pi, U_\pi)$ is an Abelian C^* -algebra, which is irreducible by assumption on π . Hence it is one-dimensional and $\pi = \pi_{x,\theta}$ where $\theta \in [0, 1)$ is given by $U_\pi \xi_x = \exp(2i\pi\theta) \xi_x$.

Now, if $x \notin \{-1, 1\}$ then let us define the linear operator $W : l^2(\mathbb{Z}) \rightarrow \mathcal{M}$ by setting $W(e_n) = U_\pi^n \xi_x$ for all $n \in \mathbb{Z}$ and where $(e_n)_{n \in \mathbb{Z}}$ is the canonical basis of $l^2(\mathbb{Z})$. Then $W^* \pi W = \pi_x$ by construction (since $W^* \pi W(A_\varphi)(e_n) = W^* A_\pi U_\pi^n \xi_x = \varphi^n(x) W^* \xi_x = \varphi^n(x) e_n$ and $W^* \pi W(U_\pi)(e_n) = W^* U_\pi U_\pi^n \xi_x = W^* U_\pi^{n+1} \xi_x = e_{n+1}$). Now, W is a unitary from $l^2(\mathbb{Z})$ onto $\mathcal{M}$ since for all $n, m \in \mathbb{Z}$ we have $\varphi^n(x) \neq \varphi^m(x)$ for $n \neq m$, hence $(U_\pi^n \xi_x)_{n \in \mathbb{Z}}$ is an orthonormal family. Hence, π is unitarily equivalent to π_x . $\square$

3.2. The Spectrum. The spectrum $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$ of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ can be identified with the set IrrRep by Theorem (3.10). We now describe the topology on $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$. Since all the representations of $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$ are of type I, the calculation of the spectrum could also use [17, Theorem 5.3].

We recall:

Proposition 3.11 ([4, Theorem VIII.2.1, p. 222]). *Given any crossed product $\mathcal{A} \rtimes_f \mathbb{Z}$ for some unital C^* -algebra $\mathcal{A}$ and f any $*$ -automorphism of $\mathcal{A}$, there exists a unique action α of the Lie group $\mathbb{T}$ on $\mathcal{A} \rtimes_f \mathbb{Z}$ which satisfies $\alpha_\lambda(U_f) = \lambda U_f$ and $\alpha_\lambda(h) = h$ for all $\lambda \in \mathbb{T}$, $h \in \mathcal{A}$ and where U_f is the canonical unitary in $\mathcal{A} \rtimes_f \mathbb{Z}$.*

Definition 3.12. *For any $a \in \mathcal{A} \rtimes_f \mathbb{Z}$ and $n \in \mathbb{Z}$ we set*

$$\mathbb{E}_n^\alpha(a) = \int_{\mathbb{T}} \alpha_\lambda(a U_f^{*n}) d\mu(\lambda)$$

where μ is the Haar probability measure on $\mathbb{T}$.

Proposition 3.13 ([4, Theorem VIII.2.2, p. 223]). *The map $\mathbb{E}_n^\alpha : \mathcal{A} \rtimes_f \mathbb{Z} \longrightarrow \mathcal{A}$ is a surjection of norm 1 for all $n \in \mathbb{Z}$, and moreover any $a \in \mathcal{A} \rtimes_f \mathbb{Z}$ is the limit of $\left(\Sigma_k^\alpha(a) = \sum_{n=-k}^k \left(1 - \frac{|n|}{k+1}\right) \mathbb{E}_n^\alpha(a) U^n \right)_{k \in \mathbb{N}}$.*

We identify the Gelfan'd spectrum $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$ of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ with the set IrrRep by Theorem (3.10):

Definition 3.14. *Let $q : \text{IrrRep} \longrightarrow \mathcal{C} \cup (\mathbb{T} \times \{-1, 1\})$, where $\mathcal{C}$ is the orbit space of $(C(\mathbb{D} \setminus \{-1, 1\}), \varphi, \mathbb{Z})$, be defined by: $q(\pi_x) = \mathcal{O}_\varphi(x) \in \mathcal{C}$ and $q(\pi_{\theta, \varepsilon}) = (e^{2i\pi\theta}, \varepsilon) \in \mathbb{T} \times \{-1, 1\}$*

Definition 3.15. *We endow $\mathcal{C} \cup (\mathbb{T} \times \{-1, 1\})$ with the topology $\mathfrak{C}$ where a set F is closed if and only if it is either contained and closed in $\mathbb{T} \times \{-1, 1\}$, or it is of the form $F' \cup (\mathbb{T} \times \{-1, 1\})$ where F' is closed in $\mathcal{C}$.*

Theorem 3.16. *Let φ be an hyperbolic automorphism of the closed unit disk $\mathbb{D}$. The map q is an homeomorphism of the spectrum $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$ of the C^* -crossed product $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ with $(\mathcal{C} \cup (\mathbb{T} \times \{-1, 1\}), \mathfrak{C})$.*

Proof. We already know that $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$ is identified, as a set, with IrrRep by Theorem (3.10). Since $\mathbb{D} \setminus \{-1, 1\}$ is an open φ -invariant set, we have the exact sequence

$$0 \rightarrow C(\mathbb{D} \setminus \{-1, 1\}) \rtimes_\varphi \mathbb{Z} \rightarrow C(\mathbb{D}) \rtimes_\varphi \mathbb{Z} \rightarrow C(\{-1, 1\}) \rtimes_\varphi \mathbb{Z} \rightarrow 0.$$

Let IrrRep_1 be the subset of IrrRep consisting of the one-dimensional representations $\pi_{\theta, \varepsilon}$ ($\varepsilon \in \{-1, 1\}, \theta \in [0, 1)$), and let IrrRep_∞ be the subset of IrrRep consisting of the infinite dimensional representations π_x in IrrRep , where $x \in \mathbb{D} \setminus \{-1, 1\}$. Observe that

$$q \left(\text{IrrRep}_1 \right) = \mathbb{T} \times \{-1, 1\} \quad \text{and} \quad q \left(\text{IrrRep}_\infty \right) = \mathcal{C}.$$

An irreducible $*$ -representation π is unitarily conjugated to some element of IrrRep_1 if and only if π vanishes on the ideal $C(\mathbb{D} \setminus \{-1, 1\}) \rtimes_\varphi \mathbb{Z}$. Let $\pi_{\theta, \varepsilon} \in \text{IrrRep}_1$. Then the quotient representation of $C(\{-1, 1\} \times \mathbb{T})$ defined by $\pi_{\theta, \varepsilon}$ is obviously $f \in C(\{-1, 1\} \times \mathbb{T}) \mapsto f(\varepsilon, e^{2i\pi\theta}) = f \circ q(\pi_{\theta, \varepsilon})$. By [6, Proposition 3.2.1, p. 61], the map q is an homeomorphism onto the spectrum of $C(\{-1, 1\}) \rtimes_\varphi \mathbb{Z}$, and IrrRep_1 is closed in $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$. Hence, a subset F of IrrRep_1 is closed in $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$ if and only if $q(F)$ is closed.

Let F be a nonempty closed set in $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$. By [6, Proposition 3.1.2, p. 60], there exists a (nonempty) subset $\mathcal{F}$ of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ such that $F = \{\pi \in \text{IrrRep} : \mathcal{F} \subseteq \ker \pi\}$. Assume that $\pi_x \in F$ for some

$x \in \mathcal{D}_\varphi$. The range of π_x is $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$. We denote by χ the canonical surjection of $\mathcal{T}$ onto $\mathcal{T}/\mathcal{K} = C(\mathbb{T})$. Let $P = 1 \oplus 0$ and $Q = 1 - P$. For any $\theta \in [0, 1)$ and any $a \in C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$, we set:

$$\chi_\theta^1(a) = \chi(P\pi_x(a)P)(e^{2i\pi\theta}) \quad \text{and} \quad \chi_\theta^{-1}(a) = \chi(Q\pi_x(a)Q)(e^{2i\pi\theta}).$$

Of course, χ_θ^1 and χ_θ^{-1} are one dimensional *-representations of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ and so they are unitarily equivalent to, respectively, $\pi_{1,\theta}$ and $\pi_{-1,\theta}$. Hence if $\pi_x(a) = 0$ then $\pi_{\theta,\varepsilon}(a) = 0$ for all $\theta \in [0, 1)$ and $\varepsilon \in \{-1, 1\}$. Hence, for all $a \in \mathcal{F}$ we have $\pi'(a) = 0$ for all π for all $\pi' \in \text{IrrRep}_1$. Thus F contains IrrRep_1 . Now, let $\pi \in F \cap \text{IrrRep}_\infty$: by Theorem (3.10), there exists $x \in \mathcal{D}_\varphi$ such that $\pi = \pi_x$. We set:

$$\mathcal{E} = \{y \in \mathbb{D} \setminus \{-1, 1\} : \exists n \in \mathbb{Z} \quad \pi_{\varphi^n(y)} \in F\}.$$

Denote by α the canonical action of $\mathbb{T}$ on $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ and β the canonical action of $\mathbb{T}$ on $C(\overline{\mathbb{D}}) \rtimes_\varphi \mathbb{Z}$. We then have, for all $\lambda \in \mathbb{T}$, that $\pi_x \circ \alpha_\lambda = \beta_\lambda \circ \pi_x$ (as can be checked on A_φ and U_φ and then extended to $C^*(A_\varphi, U_\varphi)$ since $\pi_x \circ \alpha$ and $\beta \circ \pi_x$ are *-morphisms). For all $a \in \mathcal{F}$ we have $\pi_y(a) = 0$, so $\mathbb{E}_n^\alpha(a)(y) = \pi_y(\mathbb{E}_n^\alpha(a)) = \int_{\mathbb{T}} \beta_\lambda \circ \pi_y(a U_\varphi^{*n}) d\mu(\lambda) = 0$ for all $y \in \mathcal{E}$ and for all $n \in \mathbb{Z}$. Thus, if we set:

$$\mathcal{F}' = \{a \in A : \forall n \in \mathbb{Z} \quad \forall y \in \mathcal{E} \quad \mathbb{E}_n(a)(y) = 0\}$$

then $\{\pi \in \text{IrrRep} : \mathcal{F}' \subseteq \ker \pi\} \subseteq F$. Conversely, let $a \in C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ such that $\mathbb{E}_n^\alpha(a)(y) = 0$ for all $y \in \mathcal{E}, n \in \mathbb{Z}$. Then $\Sigma_k^\alpha(a)(y) = \pi_y(\Sigma_k^\alpha(a)) = 0$ for all $y \in \mathcal{E}, k \in \mathbb{N}$, and thus $\pi_y(a) = 0$ for all $y \in \mathcal{E}$ as π_y continuous and $(\Sigma_k^\alpha(a))_{k \in \mathbb{N}}$ converges in norm to a . Hence $\{\pi \in \text{IrrRep} : \mathcal{F}' \subseteq \ker \pi\} = F$. Now, let $(\omega_k)_{k \in \mathbb{N}}$ be a sequence in $\mathcal{C}$ such that $\omega_k \in q(F \cap \text{IrrRep}_\infty)$ for all $k \in \mathbb{N}$ and converging to $\omega \in \mathcal{C}$. Let $a \in \mathcal{F}'$. Then for any $n \in \mathbb{Z}$ and any $y \in \omega$ we have $\mathbb{E}_n^\alpha(a)(y) = 0$ by continuity. Hence, $\omega \in q(F \cap \text{IrrRep}_\infty)$, so $q(F \cap \text{IrrRep}_\infty)$ is a closed set in $\mathcal{C}$. Therefore, $q(F) = \overline{\mathbb{T} \cup q(F \cap \text{IrrRep}_\infty)}$ is closed in $\mathcal{C}$.

Conversely, let F be a set in $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$ such that $F \cap \text{IrrRep}_\infty \neq \emptyset$ and $q(F)$ is closed. Then $T = q(F \cap \text{IrrRep}_\infty)$ is closed in $\mathcal{C}$. Let:

$$\mathcal{M} = \{a \in C(\mathbb{D}) \rtimes_\varphi \mathbb{Z} : \forall n \in \mathbb{Z} \quad \mathbb{E}_n(a)(T) = \{0\}\}.$$

Let $\pi \in \text{IrrRep}_\infty$ such that $\pi(\mathcal{M}) = \{0\}$ and let $\mathcal{O} = q(\pi)$. Since $\mathbb{E}_0^\alpha$ is a surjection, for any $f \in C(\mathcal{C})$ such that $f(T) = \{0\}$ there exists $a \in \mathcal{M}$ such that $\mathbb{E}_0^\alpha(a) = f$. Therefore, for all $f \in C(\mathcal{C})$ such that $f(T) = \{0\}$ we have $f(\mathcal{O}) = 0$, so $\mathcal{O} \in \overline{T} = T$. Hence $\pi \in F$. Hence, F is closed in $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$. $\square$

A corollary of Theorem (3.16) is that the spectrum of the ideal $\mathcal{I} = C(\mathbb{D} \setminus \{-1, 1\}) \rtimes_{\varphi} \mathbb{Z}$ is the set IrrRep_{∞} of infinite dimensional representations in IrrRep , with the topology of $\mathcal{C}$. Since all the representations in IrrRep_{∞} acts on the same Hilbert space $l^2(\mathbb{Z})$, we deduce moreover that $\mathcal{I}$ is a homogenous C^* -algebra. Since $\mathcal{I}$ has continuous trace, and since $H^3(\mathcal{C}, \mathbb{Z}) = \{0\}$, we deduce by [6, Corollary 10.9.6, p. 219] that $\mathcal{I} = C(\mathcal{C}, \mathcal{K})$. Moreover, it is easy to generalize the part of the proof of Theorem (3.16) concerning $\mathcal{I}$ to any C^* -crossed product $C(X) \rtimes_f \mathbb{Z}$ where the action of $\mathbb{Z}$ on X is proper and free (so that $\mathcal{O}_{\varphi}(x)$ is homeomorphic to $\mathbb{Z}$ as a subset of X , for all $x \in X$): thus the spectrum of such crossed-products is simply the orbit space of the action with its natural quotient topology. We will provide a self-contained proof that $\mathcal{I} = C(\mathcal{C}, \mathcal{K})$ in Proposition (3.22), which uses a similar but simplified method compared to [6, Chapter 10].

Remark 3.17. *Let $x \in \mathbb{D} \setminus \{-1, 1\}$. Then by [6, Proposition 3.2.1, p. 61], we deduce from the exact sequence:*

$$0 \rightarrow \ker \pi_x \rightarrow C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z} \xrightarrow{\pi_x} \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix} \rightarrow 0$$

that the spectrum of $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$ is $\text{IrrRep}_1 \cup \{\pi_x\}$, topologized as a subset of $(\text{IrrRep}, \mathfrak{C})$. This result can be proven directly by using the exact sequence $0 \rightarrow \mathcal{K} \rightarrow \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix} \rightarrow C(\mathbb{T}) \oplus C(\mathbb{T}) \rightarrow 0$ and a method similar yet simpler than the one used in Theorem (3.16).

3.3. The Crossed-Product. The following C^* -algebra is our candidate for the full crossed-product C^* -algebra $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$:

Definition 3.18. *We denote by $\mathcal{A}$ the C^* -algebra:*

$$\left\{ \begin{array}{l} f \in C\left(\mathbb{D} \setminus \{-1, 1\}, \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}\right) \text{ such that:} \\ U^* f U = f \circ \varphi \text{ and } \forall t \in \mathbb{D} \setminus \{-1, 1\} \quad f(t) - f(0) \in \mathcal{K} \end{array} \right\}$$

where U is the bilateral unitary shift defined by $U(\zeta_n)_{n \in \mathbb{Z}} = (\zeta_{n-1})_{n \in \mathbb{Z}}$ for all $(\zeta_n)_{n \in \mathbb{Z}} \in \mathbb{H} = l^2(\mathbb{Z})$.

The norm on $\mathcal{A}$ is $\|\cdot\|_{\mathcal{A}} = \sup_{t \in \mathbb{D} \setminus \{-1, 1\}} \|\cdot\|_{C(\overline{\mathbb{Z}}) \rtimes_{\tau} \mathbb{Z}}$, which is indeed a norm (and then easily checked to be a C^* -norm) thanks to the condition $U^* f U = f \circ \varphi$, and is well-defined since $C(\overline{\mathbb{Z}}) \rtimes_{\tau} \mathbb{Z} = \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$ by Lemma (3.7).

Remark 3.19. The C^* -algebra $\mathcal{A}$ could alternatively be defined as the set of $f \in C\left(\mathbb{D} \setminus \{-1, 1\}, \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}\right)$ such that $U^*fU = f \circ \varphi$ and

$$(\chi \oplus \chi)(f(t)) = (\chi \oplus \chi)(f(0))$$

for all $t \in \mathbb{D} \setminus \{-1, 1\}$ where $\chi : \mathcal{T} \rightarrow C(\mathbb{T})$ is the canonical projection of the Toeplitz algebra on $C(\mathbb{T})$ (i.e. the symbol map). This second condition reflects the fact that φ admits two and only two distinct fixed points on $\mathbb{D}$.

Lemma 3.20. Let $A'(t) = \text{Diag}((\varphi^n(t))_{n \in \mathbb{Z}})$ acting on $\mathbb{H} = l^2(\mathbb{Z})$ for all $t \in \mathbb{D} \setminus \{-1, 1\}$. Then A' is a continuous function from $\mathbb{D} \setminus \{-1, 1\}$ into the bounded operators on $\mathbb{H}$. Let $U' : t \in \mathbb{D} \setminus \{-1, 1\} \mapsto U$ where $U \in \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$, as defined in Definition (3.4) and Definition (3.18), is the bilateral shift on $\mathbb{H}$. Then $C^*(A', U') = \mathcal{A}$.

Proof. By definition $C^*(A', U') \subseteq \mathcal{A}$: indeed, $A'(t) - A'(0) = 0 \oplus 0$ and $U'(t) - U'(0) = 0 \oplus 0$ are independent of $t \in \mathbb{D} \setminus \{-1, 1\}$. Moreover, $A \circ \varphi = U^*A'U$ and $U' \circ \varphi = U' = U^*U'U$. Let us now prove the converse inclusion.

Let $(e_n)_{n \in \mathbb{Z}}$ be the canonical basis of $l^2(\mathbb{Z})$ and let E_n be the orthogonal projection in $l^2(\mathbb{Z})$ on $\mathbb{C}e_n$. Let $\mathcal{C}_0$ be the C^* -algebra consisting of the operators $g(A')$ for $g \in C(\mathbb{D})$ such that $g(-1) = g(1) = 0$, and let $\mathcal{A}_0 = \{f \in C(\mathbb{D} \setminus \{-1, 1\}, \mathcal{K}) : U^*fU = f \circ \varphi\}$.

First, we observe that $\mathcal{C}_0$ is the C^* -subalgebra $\mathcal{C}'_0$ of $\mathcal{A}_0$ of the continuous functions on $\mathbb{D} \setminus \{-1, 1\}$ subject to the condition $U^*fU = f \circ \varphi$ and such that $f(t)$ is diagonal in $(e_n)_{n \in \mathbb{Z}}$ for all $t \in \mathbb{D} \setminus \{-1, 1\}$. First, by construction $\mathcal{C}_0 \subseteq \mathcal{C}'_0$. On the other hand, let $f \in \mathcal{C}'_0$. For each $t \in \mathbb{D} \setminus \{-1, 1\}$ we have $f(t) \in C^*(A'(t))$, so $f(t) = g_t(A'(t))$ for some continuous function $g_t \in C(\overline{\mathcal{O}_\varphi(t)})$ such that $g_t(1) = g_t(-1) = 0$ (as $f(t)$ is compact). Since f is continuous, we deduce that the function $g : t \in \mathbb{D} \setminus \{-1, 1\} \mapsto g_t(t)$ is also continuous, and moreover it can be continuously extended to $\mathbb{D}$ by setting $g(1) = g(-1) = 0$. Now, since $U^*A'(t)U = A' \circ \varphi(t)$ for all $t \in \mathbb{D} \setminus \{-1, 1\}$, it follows that $g_t(\varphi(t)) = g_{\varphi(t)}(\varphi(t))$. Hence:

$$\begin{aligned} g(A'(t)) &= g(\text{Diag}(\varphi^n(t))_{n \in \mathbb{Z}}) = \text{Diag}((g \circ \varphi^n(t))_{n \in \mathbb{Z}}) \\ &= \text{Diag}\left((g_{\varphi^n(t)}(\varphi^n(t)))_{n \in \mathbb{Z}}\right) = \text{Diag}((g_t(\varphi^n(t)))_{n \in \mathbb{Z}}) \\ &= g_t(A'(t)) = f(t). \end{aligned}$$

Hence, $f \in \mathcal{C}_0$, so indeed $\mathcal{C}_0 = \mathcal{C}'_0$.

Now, let $f \in \mathcal{A}_0$. We write $f = f_1 + f_2$ where $f_1 \in \mathcal{C}_0$ such that $f_1(iy) = f(iy)$ for all $y \in [-1, 1]$. Thus, $f_2 \in \mathcal{A}_0$ and moreover, $f_2(iy) = 0$ for all $y \in [-1, 1]$. Hence, since $U^*f_2U = f_2 \circ \varphi$, we also have $f_2(\varphi(iy)) = 0$. Thus, $f_2 \in \{g \in \mathcal{A}_0 : g(L \cup \varphi(L)) = \{0\}\}$ where $L = \{iy : y \in [-1, 1]\}$. But it is now immediate to check that $\{g \in \mathcal{A}_0 : g(L \cup \varphi(L)) = \{0\}\}$ is $*$ -isomorphic to the C^* -subalgebra of operators $g \in C(\overline{\mathcal{D}_\varphi}) \otimes \mathcal{K}$ such that $g(L \cup \varphi(L)) = \{0\}$. Now, given any function $g \in C(\overline{\mathcal{D}_\varphi})$, we define $g' : t \in \mathbb{D} \setminus \{-1, 1\} \mapsto g([t])e_{n(t)}$ where $[t]$ and $n(t)$ are the unique element in $\mathcal{D}_\varphi$ and $\mathbb{Z}$ respectively such that $\varphi^{n(t)}([t]) = t$. Now, it is easy to check that g' is continuous on $\mathbb{D} \setminus \{-1, 1\}$ (as the only problems are at the boundary, on which g is zero everywhere), and $U^*g'U = g' \circ \varphi$ as well as $g(t) \in \mathcal{K}$ and is diagonal in $(e_n)_{n \in \mathbb{Z}}$ for all $t \in \mathbb{D} \setminus \{-1, 1\}$. Hence $g' \in \mathcal{C}_0$. From this it is easy to check that $\{g \in \mathcal{A}_0 : g(L \cup \varphi(L)) = \{0\}\} \subseteq C^*(\mathcal{C}_0, U') = C^*(A', U')$. Therefore, $f_2 \in C^*(A', U')$. Hence $f \in C^*(A', U')$ and thus $\mathcal{A}_0 \subseteq C^*(A', U')$.

Now, we conclude our theorem as follows. Let $f \in \mathcal{A}$. There exists $g \in C^*(A', U')$ such that $g(0) = f(0)$ by Lemma (3.6), which proves that both $f(t)$ and $g(t)$ lie in $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$ for all $t \in \mathbb{D} \setminus \{-1, 1\}$. Now, $f(t) - g(t) \in \mathcal{K}$ so, by construction, $f - g \in \mathcal{A}_0$. Thus $f - g \in C^*(A', U')$. This shows that $f = (f - g) + g \in C^*(A', U')$.

Hence as claimed, $\mathcal{A} = C^*(A', U')$. $\square$

Theorem 3.21. *Let φ be a hyperbolic automorphism of the closed unit disk $\mathbb{D}$. Then $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ is $*$ -isomorphic to:*

$$\mathcal{A}' = \left\{ \begin{array}{l} f \in C\left(\mathbb{R} \times [0, 1], \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}\right) \text{ such that:} \\ \forall x \in \mathbb{R} \ \forall y \in [0, 1] \ \left\{ \begin{array}{l} f(x, y) - f(0, 0) \in \mathcal{K} \\ U^*f(x, y)U = f(x + 1, y) \end{array} \right. \end{array} \right\}.$$

Proof. First we observe that $\sigma(A') = \mathbb{D}$ and $U'^*A'U' = \varphi(A')$. Hence, there exists a unique $*$ -epimorphism θ from $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ onto $C^*(A', U')$ such that $\theta(A_\varphi) = A'$ and $\theta(U_\varphi) = U'$. Let $a \in C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ such that $\theta(a) = 0$. Let π be an irreducible $*$ -representation of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$. By Theorem (3.10), up to unitary equivalence, $\pi \in \text{IrrRep}$. Let $x \in \mathbb{D}$ such that $\sigma(\pi(A_\varphi)) = \overline{\mathcal{O}_\varphi(x)}$. If $x \in \mathbb{D} \setminus \{-1, 1\}$ then π acts on $l^2(\mathbb{Z})$ and extends: $\pi(A_\varphi)f(z) = \varphi(x)^z f(z)$ and $\pi(U_\varphi)f(z) = f(z - 1)$. We set ρ to be the faithful $*$ -representation of $\begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}$ on $l^2(\mathbb{Z})$ given in Lemma (3.6). Let $\pi' : f \in \mathcal{A} \mapsto \rho(f(x))$: by construction, π' is an (irreducible) $*$ -representation of $C^*(A', U')$.

If $x = \{-1, 1\}$ then π acts on $\mathbb{C}$ by $\pi(U_\varphi) = \lambda \in \mathbb{T}$ and $\pi(A_\varphi) = x$. If $x = 1$ then we set $\pi' : f \in \mathcal{A} \mapsto ((0 \oplus \chi) f(t))(\lambda)$ (for any $t \in \mathbb{D} \setminus \{-1, 1\}$ since $(0 \oplus \chi)(f(t))$ is independent of t for $f \in \mathcal{A}$ by construction). If $x = -1$ we set $\pi' : f \in \mathcal{A} \mapsto ((\chi \oplus 0) f(t))(\lambda)$.

Whatever $x \in \mathbb{D}$ is, it is now easy to check that $\pi' \circ \theta = \pi$, as this equality is valid on the total set of monomials in A_φ, U_φ . Therefore, $\pi(a) = \pi' \circ \theta(a) = 0$. Since π is arbitrary, we deduce $a = 0$. Hence θ is also injective, hence θ is a *-isomorphism from $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ onto:

$$\mathcal{A} = \left\{ \begin{array}{l} f \in C\left(\mathbb{D} \setminus \{-1, 1\}, \left[\begin{array}{cc} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{array} \right] \right) \text{ such that:} \\ U^* f U = f \circ \varphi \text{ and } \forall t \in \mathbb{D} \setminus \{-1, 1\} \quad f(t) - f(0) \in \mathcal{K} \end{array} \right\}.$$

Now, in the proof of Theorem (2.4), we constructed an homeomorphism μ from $\mathcal{D}_\varphi$ onto $[0, 1]^2$, which can be extended by induction to an homeomorphism from $\mathbb{D} \setminus \{-1, 1\}$ onto $\mathbb{R} \times [0, 1]$ in such a way that $\mu^{-1} \circ \varphi \circ \mu$ is simply the translation $(x, y) \in \mathbb{R} \times [0, 1] \mapsto (x + 1, y)$. The map $\psi : f \in \mathcal{A} \mapsto f \circ \mu \in \mathcal{A}'$ is obviously a *-isomorphism. $\square$

Since the action of $\mathbb{Z}$ on $\mathbb{D} \setminus \{-1, 1\}$ by an hyperbolic automorphism fixing $\{-1, 1\}$ is proper and free, the ideal $C(\mathbb{D} \setminus \{-1, 1\}) \rtimes_\varphi \mathbb{Z}$ of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ is $C(\mathcal{C}, \mathcal{K})$, as can be derived from our work or the general theory of such crossed-products (which are known to be field of elementary C*-algebras over the orbit space, thus classified by their Dixmier -Douady class, which lives in $H^3(\mathcal{C}, \mathbb{Z}) = \{0\}$, hence the field is trivial). We offer here the proof of this last isomorphism as a corollary of our work:

Proposition 3.22. *Let φ be a hyperbolic automorphism of $\mathbb{D}$. Then $C_{-1,1}(\mathbb{D}) \rtimes_\varphi \mathbb{Z} = C(\mathcal{C}, \mathcal{K})$ where $\mathcal{C} = [0, 1] \times \mathbb{T}$ and $C_{-1,1}(\mathbb{D})$ is the C*-algebra of continuous functions on $\mathbb{D}$ such that $f(-1) = f(1) = 0$.*

Proof. The C*-algebra $C_{-1,1}(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ is the kernel of the map η by construction, so it is isomorphic to $C(\mathbb{D} \setminus \{-1, 1\}, \mathcal{K} : U^* f U = f \circ \varphi)$. There exists a selfadjoint operator H on $l^2(\mathbb{Z})$ such that $U = \exp(iH)$. Let $U_t = \exp(itH)$ for all $t \in [0, 1]$. Let h be an homeomorphism from $\mathbb{D} \setminus \{-1, 1\} \longrightarrow \mathbb{R} \times [0, 1]$ such that $h\varphi h^{-1}$ is just the translation by 1 on $\mathbb{R}$. For any $f \in C(\mathcal{C}, \mathcal{K})$ we define $\tilde{f}$ as follows: $\tilde{f}(x) = U_r^* f(e(r_0), s) U_r$ where $h(r, s) = \omega$. Now, $\tilde{f} \in C(\mathbb{D} \setminus \{-1, 1\}, \mathcal{K} : U^* f U = f \circ \varphi)$. The converse inclusion is shown similarly. $\square$

3.4. K-Theory, Projections and Unitaries.

3.4.1. *K-theory for general actions on $\mathbb{D}$.* The whole K -theory of $\mathbb{D}$ can be read from the following six-terms exact sequence [2, Theorem 9.3.1, p. 67]:

$$\begin{array}{ccccc} K_0\left(\overset{\circ}{\mathbb{D}}\right) = \mathbb{Z} & \longrightarrow & K_0(\mathbb{D}) = \mathbb{Z} & \longrightarrow & K_0(\mathbb{T}) = \mathbb{Z} \\ \uparrow & & & & \downarrow \\ K_1(\mathbb{T}) = \mathbb{Z} & \longleftarrow & K_1(\mathbb{D}) & \longleftarrow & K_1\left(\overset{\circ}{\mathbb{D}}\right) = 0 \end{array}$$

which follows from the exact sequence

$$0 \longrightarrow C\left(\overset{\circ}{\mathbb{D}}\right) \longrightarrow C(\mathbb{D}) \longrightarrow C(\mathbb{T}) \longrightarrow 0.$$

We conclude immediately that $K_0(\mathbb{D}) = \mathbb{Z}$ and $K_1(\mathbb{D}) = 0$. Yet, since $\mathbb{D}$ is a contractible compact metric space, we immediately conclude that $K_0(\mathbb{D}) = \mathbb{Z}$ with generator [1]. From this, we can deduce the K -theory of the crossed-product of φ on $C(\mathbb{D})$:

Proposition 3.23. *We have $K_0(C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}) = \mathbb{Z}$, generated by the class of the identity, and $K_1(C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}) = \mathbb{Z}$ generated by the canonical unitary of $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$, for any homeomorphism φ of $\mathbb{D}$.*

Proof. The Pimsner-Voiculescu six-terms exact sequence [13], [2, Theorem 10.2.1, p. 73] for this action is:

$$\begin{array}{ccccc} K_0(\mathbb{D}) = \mathbb{Z} & \xrightarrow{1-\varphi^*} & K_0(\mathbb{D}) = \mathbb{Z} & \longrightarrow & K_0(C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}) \\ \uparrow & & & & \downarrow \\ K_1(C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}) & \longleftarrow & K_1(\mathbb{D}) = 0 & \xleftarrow{1-\varphi^*} & K_1(\mathbb{D}) = 0 \end{array}$$

and deduce the result immediately since $(1 - \varphi^*)[1] = 0$. Note that the K -groups do not depend on the homeomorphism φ of $\mathbb{D}$. $\square$

3.4.2. *Another view of K-theory.* Even though the only nontrivial class in $K_0(C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z})$ is always the one of the identity, there can be many nontrivial projections in $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ whose K_0 classes vanish. This is illustrated by our example, when φ is an hyperbolic automorphism:

Proposition 3.24. *Let φ be a hyperbolic automorphism of $\mathbb{D}$. The projections of $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ of K -class 0 are (in one-to-one correspondence with) the projections in $C(\mathcal{C}, \mathcal{K})$. Any projection P of $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ is homotopic to Q or $1-Q$ for some compact-projection-valued continuous function Q on $\mathbb{D} \setminus \{-1, 1\}$ such that $U^*QU = Q \circ \varphi$.*

The unitaries of $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ are all homotopic to U^n for some $n \in \mathbb{Z}$.

Proof. By Proposition (3.22) we have the following exact sequence:

$$0 \longrightarrow C(\mathcal{C}, \mathcal{K}) \longrightarrow C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z} \longrightarrow C(\mathbb{T}) \oplus C(\mathbb{T}) \longrightarrow 0$$

which implies that the following six-terms exact sequence holds:

$$\begin{array}{ccccccc} K_0(\mathcal{C}, \mathcal{K}) = \mathbb{Z} & \longrightarrow & K_0(C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}) = \mathbb{Z} & \longrightarrow & \mathbb{Z}^2 & & \\ \uparrow & & & & \downarrow & & \\ \mathbb{Z}^2 & \longleftarrow & K_1(C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}) = \mathbb{Z} & \longleftarrow & K_1(\mathcal{C}, \mathcal{K}) = \mathbb{Z} & & \end{array}.$$

By inspecting each arrow, we conclude that the image of $K_0(\mathcal{C}, \mathcal{K})$ can only be $\{0\}$ for the top row to be exact. Thus all compact projections in $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ have null K_0 -class. Since obviously $1 \in C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ has for image $(1, 1) \in \mathbb{Z}^2$, we conclude that there is no projections in $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ which maps to either $0 \oplus n$ or $n \oplus 0$ for any $n \neq 0$. Thus for any projection $P \in C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ there exists a projection $K \in C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ with $K(t)$ compact for all $t \in \mathbb{D} \setminus \{-1, 1\}$ such that P is homotopic (in the space of projections) to K or $1 - K$.

Similarly, the bottom row exactness imposes that $K_1(C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z})$ maps into a copy of $\mathbb{Z}$ in $\mathbb{Z}^2$ and since U has class $(1, -1)$ (since the image of U in $C(\mathbb{T}) \oplus C(\mathbb{T})$ is by construction $z \mapsto z \oplus \bar{z}$), we conclude that any unitary W in $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ which K_1 -class $n \in \mathbb{Z}$ is of the form $V + K$ where $K \in C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ is compact-valued (i.e. $\eta(K) = 0$) and V is homotopic (in the unitary group) to U^n . $\square$

In particular, one observes that if $\mathcal{A}'$ is the C^* -algebra:

$$\left\{ f \in C\left(\mathcal{C}, \begin{bmatrix} \mathcal{T} & \mathcal{K} \\ \mathcal{K} & \mathcal{T} \end{bmatrix}\right) : (\chi \oplus \chi)(f(t) - f(0)) = 0 \right\}$$

then $K_0(\mathcal{A}') = \mathbb{Z}^2$, so Proposition (3.22) does not extend to $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$.

4. PARABOLIC AUTOMORPHISMS

Informally, a parabolic automorphism of the closed unit disk $\mathbb{D}$ is a limit case for the hyperbolic automorphisms we studied in the previous section. However, the situation turns out to be easier in the parabolic case. Let φ be a parabolic automorphism of $\mathbb{D}$ which, by Theorem (2.4), we can assume to fix 1.

First, as in the hyperbolic case, there exists a domain $\mathcal{D}_{\varphi}$ such that for all $x \in \mathbb{D} \setminus \{1\}$ there exists a unique $y \in \mathcal{D}_{\varphi}$ such that $\mathcal{O}_{\varphi}(x) = \mathcal{O}_{\varphi}(y)$. For instance, one can choose $\mathcal{D}_{\varphi}$ as the region between the diameter passing by 1 and its image by φ , not including this image. Second, observe that the orbit space of φ on $\mathbb{D}$ is the compact cone $\mathfrak{S} = \{(t, t\omega) : t \in [0, 1], \omega \in \mathbb{T}\}$, which is homeomorphic to $\mathbb{D}$.

Definition 4.1. Let φ be a parabolic automorphism of $\mathbb{D}$ with fixed point 1 (necessarily unique).

Let $\theta \in [0, 1)$. We define the $*$ -representation π_θ of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ on $\mathbb{C}$ by setting:

$$\pi_\theta(A_\varphi) = 1 \quad \text{and} \quad \pi_\theta(U_\varphi) = \exp(2i\pi\theta).$$

Let now $x \in \mathbb{D} \setminus \{1\}$. Then we define the $*$ -representation π_x of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ on $l^2(\mathbb{Z})$ by setting:

$$\pi_x(A_\varphi)((\xi_n)_{n \in \mathbb{Z}}) = (\varphi^n(x)\xi_n)_{n \in \mathbb{Z}} \quad \text{and} \quad \pi_x(U_\varphi)((\xi_n)_{n \in \mathbb{Z}}) = ((\xi_{n-1})_{n \in \mathbb{Z}})$$

for all $(\xi_n)_{n \in \mathbb{Z}} \in l^2(\mathbb{Z})$.

We set $\text{IrrRep} = \{\pi_\theta, \pi_x : \theta \in [0, 1), x \in \mathcal{D}_\varphi\}$.

Theorem 4.2. Let φ be a parabolic conformal automorphism of $\mathbb{D}$. Then all the representations in IrrRep are irreducible $*$ -representations, and no two representations in IrrRep are unitarily equivalent. Moreover, let π be an irreducible representation of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$. Then π is unitarily equivalent to a representation in IrrRep .

Proof. It is straightforward to check that IrrRep is indeed a set of representations. If $\pi, \pi' \in \text{IrrRep}$ and $\pi \neq \pi'$ then $\sigma(\pi(U_\varphi)) \neq \sigma(\pi'(U'_\varphi))$ or $\sigma(\pi(A_\varphi)) \neq \sigma(\pi'(A_\varphi))$ so either way π and π' are not unitarily conjugated.

Let π be an irreducible representation of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$. By Theorem (3.2) and denoting $\pi(A_\varphi)$ by A_π , we have $\sigma(A_\pi) = \overline{\mathcal{O}_\varphi(x)}$ for some $x \in \mathbb{D}$. The same proof as Theorem (3.10) now applies: $x \in \overline{\mathcal{O}_\varphi(x)}$ is isolated in $\overline{\mathcal{O}_\varphi(x)}$ for φ parabolic, so there exists a vector ξ_x of norm 1 such that $A_\pi \xi_x = x \xi_x$. By irreducibility of π , it follows that $\mathcal{H} = \overline{\text{span}}\{U^n \xi_x : n \in \mathbb{Z}\}$. If $x = 1$ then $\mathcal{H} = \mathbb{C}$ and then $\pi = \pi_\theta$ for $\theta \in [0, 1)$. If $x \neq 1$ then $(U^n \xi_x)_{n \in \mathbb{Z}}$ is orthonormal, and thus if we define the operator $W : l^2(\mathbb{Z}) \rightarrow \mathcal{M}$ by linearity such that $W(e_n) = U^n \xi_x$ for all $n \in \mathbb{Z}$, where $(e_n)_{n \in \mathbb{Z}}$ is the canonical basis for $l^2(\mathbb{Z})$, then $W^* \pi W = \pi_x$. $\square$

We now identify the range of the infinite dimensional irreducible representations of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$:

Lemma 4.3. Let $x \in \mathbb{D} \setminus \{1\}$, and denote $\pi_x(A_\varphi)$ by A_x and $\pi_x(U_\varphi)$ by U . Then $C^*(U, A_x) = C^*(U) + \mathcal{K}$ where $\mathcal{K}$ is the C^* -algebra of compact operators on $l^2(\mathbb{Z})$.

Proof. Let $x \in \mathbb{D} \setminus \{1\}$. Let $B \in \pi_x(C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}) = C^*(A_x, U)$. Hence, the operator B is the limit in norm of polynomials in A_x and U . Now, $A_x^k - 1^k I$ is a compact operator, so $U^{*n} (A_x^k - 1^k I) U^n U^{m-n}$ is a compact operator. Hence $C^*(A_x, U) \subseteq \mathcal{K} + C^*(U)$.

Now, let C be a finite-rank operator on $l^2(\mathbb{Z})$: then $C = \sum_{k=-n}^n C_k U^k$ where C_k are finite rank diagonal operators. Now, trivially $C_k = f_k(A_x)$ for some continuous function f_k on $C(\overline{\mathcal{O}(x)})$ such that $f_k(1) = 0$. Hence, $C \in C^*(A_x, U)$ and thus $\mathcal{K} + C^*(U) = C^*(A_x, U)$. $\square$

Remark 4.4. *It is easy to prove that $C^*(A_x, U)$ is the C^* -crossed-product $C(\mathbb{Z} \cup \{\infty\}) \rtimes_{\tau} \mathbb{Z}$ where $\tau(z) = z + 1$ and $\tau(\infty) = \infty$, and where $\mathbb{Z} \cup \{\infty\}$ is the one-point compactification of $\mathbb{Z}$.*

We now bundle all these irreducible representations together to form the crossed-product C^* -algebra:

Theorem 4.5. *Let φ be a parabolic conformal automorphism of the closed unit disk $\mathbb{D}$ with fixed point $1 \in \mathbb{T}$. Let $\mathbb{H} = l^2(\mathbb{Z})$ and let U be the bilateral shift $U(\xi_z)_{z \in \mathbb{Z}} = (\xi_{z-1})_{z \in \mathbb{Z}}$ for all $\xi \in \mathbb{H}$. We also define, for all $t \in \mathbb{D} \setminus \{1\}$, the operator $A_x(\xi_z)_{z \in \mathbb{Z}} = (\varphi^z(x)\xi_z)_{z \in \mathbb{Z}}$. Let $U' : t \in \mathbb{D} \setminus \{1\} \mapsto U$ and $A' : t \in \mathbb{D} \setminus \{1\} \mapsto A_t$. Then $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ is $*$ -isomorphic to the C^* -algebra $C^*(A', U')$, or equivalently to the C^* -algebra:*

$$\mathcal{A} = C^*(U') + C_1(\mathbb{D}, \mathcal{K})$$

where $C_1(\mathbb{D}, \mathcal{K})$ is the C^* -algebra of $\mathcal{K}$ -valued continuous functions on $\mathbb{D}$ vanishing at 1.

Proof. Since $A' = I + K = U'U'^* + K$ where $K \in C_1(\mathbb{D}, \mathcal{K})$, it is obvious that $C^*(A', U') \subseteq C^*(U') + C_1(\mathbb{D}, \mathcal{K})$. We can proceed as in Lemma (3.20) to prove that $C^*(A', U') = C^*(U') + C_1(\mathbb{D}, \mathcal{K})$: if $f \in C^*(U') + C_1(\mathbb{D}, \mathcal{K})$ then $f = g + h$ where $g \in C^*(U')$ and $h \in C_1(\mathbb{D}, \mathcal{K})$. Now $C_1(\mathbb{D}, \mathcal{K}) = \{h \in C(\mathbb{D}) : h(1) = 0\} \otimes \mathcal{K}$. Yet, $C^*(A')$ is the C^* -algebra of continuous functions on $\mathbb{D}$ vanishing at 1 and valued in the diagonal compact operators for the canonical basis of $l^2(\mathbb{Z})$. Thus, $C_1(\mathbb{D}, \mathcal{K}) \subseteq C^*(A', U')$ trivially, and thus $f \in C^*(A', U')$ as desired.

Therefore $\mathcal{A}$ is generated by U' and A' such that $U'^*A'U' = \varphi(A')$ and $\sigma(A') = \mathbb{D}$, thus there exists a $*$ -epimorphism $\psi : C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z} \rightarrow \mathcal{A}$ such that $\psi(A_{\varphi}) = A'$ and $\psi(U_{\varphi}) = U'$. Now, let π be an irreducible $*$ -representation of $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$. Up to unitarily equivalence, $\pi \in \text{IrrRep}$. Let $a \in C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ such that $\psi(a) = 0$. Assume first $\pi = \pi_{\theta}$ for some $\theta \in [0, 1)$. Define $\pi'(U') = e^{2i\pi\theta}$ and $\pi'(A') = 1$. Thus π' extends to a $*$ -representation of $\mathcal{A}$ and by construction, $\pi' \circ \psi = \pi$, so $\pi(a) = 0$. Assume now $\pi = \pi_x$ for $x \in \mathbb{D} \setminus \{1\}$, as defined in Lemma (4.3). Then set $\pi'(A_{\varphi}) = A_x$ and $\pi'(U_{\varphi}) = U$. Again, we have $\pi(a) = \pi' \circ \psi(a) = 0$. Hence ψ is injective, which concludes our proof. $\square$

We conclude this section with a description of $\widehat{C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}}$. We work out the spectrum from Theorem (4.5) rather than from pure representation theory considerations as in Theorem (3.16). Let $\mathbb{D}$ be the orbit space of $(C(\mathbb{D}), \varphi, \mathbb{Z})$ with its natural quotient topology.

Theorem 4.6. *Let φ be a parabolic automorphism of the closed unit disk $\mathbb{D}$. We define the map $q : \text{IrrRep} \mapsto \mathbb{T} \cup \mathbb{D} \setminus \{1\}$ by setting: $q(\pi_{\theta}) = \lambda$ and $q(\pi_x) = x$. We define on $\mathbb{T} \cup \mathbb{D} \setminus \{1\}$ the topology $\mathfrak{C}$ where a subset $F \subseteq \mathbb{T} \cup \mathbb{D} \setminus \{1\}$ is closed if and only if it is either contained and closed in $\mathbb{T}$, or it is of the form $F' \cup \mathbb{T}$ where F' is closed in $\mathbb{D} \setminus \{1\}$. The spectrum $\widehat{C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}}$ of $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ is homeomorphic to $(\mathbb{T} \cup \mathbb{D} \setminus \{1\}, \mathfrak{C})$.*

Proof. By Theorem (4.5), we have $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z} = C^*(U') + C_1(\mathbb{D}, \mathcal{K})$. For any $f \in C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ we write $f = f_U + f_{\mathcal{K}}$ where $f_U \in C^*(U')$ and $f_{\mathcal{K}} \in C_1(\mathbb{D}, \mathcal{K})$ are uniquely determined by this equation. Let π be an irreducible *-representation of $C^*(U') + C_1(\mathbb{D}, \mathcal{K})$. Then π restricted to the ideal $C_1(\mathbb{D}, \mathcal{K})$ is also irreducible, so it is either 0 or the evaluation map at some $x \in \mathbb{D} \setminus \{1\}$. In the first case, then π is really an irreducible representation of $C^*(U)$, namely $\pi = \pi_{\theta}$ for some $\theta \in [0, 1)$, and in the second case $\pi = \pi_x$. Now, we have: $\pi_{\theta}(f) = f_U(e^{2i\pi\theta})$ and $\pi_x(f) = f_U + f_{\mathcal{K}|_{\mathcal{O}(x)}}$ where $f_{\mathcal{K}|_{\mathcal{O}(x)}}$ is the restriction of $f_{\mathcal{K}}$ to $\mathcal{O}(x)$.

Let F be a subset of $\widehat{C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}}$. Let $f \in \cap_{\pi \in F} \ker \pi$. Suppose first that there exists $x \in \mathbb{D} \setminus \{1\}$ such that $\pi_x \in F$. Let y in the closure of $G = \{\mathcal{O}(x) \in \mathbb{D} : \pi_x \in F\}$ for $\mathfrak{C}$. Then either $y \in \mathbb{T}$ or $y \in \overline{G} \cap \mathcal{O}$. Now, $\pi_x(f) = f_U + f_{\mathcal{K}|_{\mathcal{O}(x)}} = 0$, so $f_U = 0$ and thus $\pi_y(f) = 0$ for all $y \in \mathbb{T}$. On the other hand, since $f_{\mathcal{K}}$ is continuous, $f_{\mathcal{K}|_y} = 0$ if y is the limit in $\mathbb{D}$ of elements in G . Thus, F is closed and contains an infinite dimensional irreducible representation if and only if $y \in \overline{G}$, hence if and only if G is closed in $\mathfrak{C}$. On the other hand if F contains only one dimensional representations, then $F = \{\pi_{\theta} : e^{2i\pi\theta} \in G\}$ for some $G \subseteq \mathbb{T}$ is closed if and only if G is closed. This concludes the description of the topology on $\widehat{C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}}$. $\square$

5. ELLIPTIC AUTOMORPHISMS

The elliptic maps naturally give rise to slight generalization of the rotation algebras. For any $\theta \in [0, 1]$ we denote by $\mathcal{A}_{\theta}$ the universal C^* -algebra generated by two unitaries U and V such that $VU = e^{2i\pi\theta}UV$ [14]. We first observe that $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ is by definition the universal C^* -algebra generated by one unitaries U_{φ} and one normal element A_{φ} such that $\sigma(A_{\varphi}) = \mathbb{D}$ and $U_{\varphi}^* A_{\varphi} U_{\varphi} = e^{2i\pi\theta} A_{\varphi}$.

Proposition 5.1. *Let $\theta \in [0, 1)$, and for any $r \in [0, 1]$ we denote by Γ_r the circle of center 0 and radius r . We let ξ_r be defined as the quotient map:*

$$C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z} \xrightarrow{\xi_r} \begin{cases} C(\Gamma_r) \rtimes_{\varphi} \mathbb{Z} = \mathcal{A}_{\theta} & \text{if } r > 0, \\ C(\Gamma_0) \rtimes_{\varphi} \mathbb{Z} = C(\mathbb{T}) & \text{if } r = 0. \end{cases}$$

Then π is an irreducible representation of $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ if and only if either of the following holds:

- *There exists $r \in]0, 1]$ and ρ an irreducible representation of $\mathcal{A}_{\theta}$ such that $\pi = \rho \circ \xi_r$,*
- *There exists $\omega \in \mathbb{T}$ such that $\pi = \nu_{\omega} \circ \xi_0$, where $\nu_{\omega}(f) = f(\omega)$ for all $f \in C(\mathbb{T})$.*

Consequently, two representations $\rho \circ \xi_r$ and $\rho' \circ \xi_{r'}$ are unitarily equivalent if and only if $r = r'$ and ρ and ρ' are unitarily equivalent representations of $C(\Gamma_r) \rtimes_{\varphi} \mathbb{Z}$.

Proof. Since $\varphi(\Gamma_r) = \Gamma_r$ for all $r \in [0, 1]$, the following exact sequence holds:

$$0 \longrightarrow C(\mathbb{D} \setminus \Gamma_r) \rtimes_{\varphi} \mathbb{Z} \longrightarrow C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z} \xrightarrow{\xi_r} C(\Gamma_r) \rtimes \mathbb{Z} \longrightarrow 0$$

so ξ_r is well-defined. Let π be an irreducible *-representation of $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$. Then by Theorem (3.2), there exists $r \in [0, 1]$ such that $\sigma(\pi(A_{\varphi})) \subseteq \Gamma_r$. It follows naturally that $\pi(C(\mathbb{D} \setminus \Gamma_r) \rtimes_{\varphi} \mathbb{Z}) = \{0\}$ and thus there exists a *-representation ρ of $C(\Gamma_r) \rtimes \mathbb{Z}$ such that $\pi = \rho \circ \xi_r$. Since π is irreducible, so is ρ . If $r > 0$ then $C(\Gamma_r) \rtimes \mathbb{Z} = \mathcal{A}_{\theta}$, while if $r = 0$ then $C(\Gamma_0) \rtimes \mathbb{Z} = C(\mathbb{T})$, and this concludes the necessary condition. Now, the conditions are trivially sufficient, and the unitary conjugacy classification follows immediately. $\square$

We now provide the description of the crossed-product C^* -algebra for elliptic automorphisms, starting by the following lemma:

Lemma 5.2. *Let $A' : t \in [0, 1] \mapsto tV \in \mathcal{A}_{\theta}$ and $U' : t \in [0, 1] \mapsto U \in \mathcal{A}_{\theta}$. Then $C^*(A', U') = \{f \in C([0, 1], \mathcal{A}_{\theta}) : f(0) \in C^*(U)\}$.*

Proof. Let $f \in C([0, 1], \mathcal{A}_{\theta})$ such that $f(0) \in C^*(U)$. We write $f = f_1 + f_2$ where $f_1 : t \in [0, 1] \mapsto f(0) \in C^*(U)$. Obviously, $f_1 \in C^*(U')$. Now, $f_2 \in C_0([0, 1]) \otimes \mathcal{A}_{\theta}$. Observe that $t \in [0, 1] \mapsto t^2 = A'^* A'(t) \in C^*(A', U')$. By the Stone-Weierstrass theorem, we deduce that $C^*(A'^* A') = C_0([0, 1])$. Let $\varepsilon > 0$ and $m \in \mathbb{Z}$, and let $\delta > 0$ such that $g(t) \leq \varepsilon$ for $t \in [0, \delta]$. Let $h_{\delta} \in C_0([0, 1])$ such that $h_{\delta}(t) = t^{-m}$ if $t \geq \delta$, while $h_{\delta}(t) \leq \frac{1}{t^m}$ if $t \in [0, \delta]$ with $h_{\delta}(0) = 0$. Then $\|g(t)V^m - g(t)h_{\delta}(t)t^m V^m\| = 0$ if $t \geq \delta$ and if $t \in [0, \delta]$ then $\|g(t)V^m - g(t)h_{\delta}(t)t^m V^m\| \leq |g(t)| |1 - h_{\delta}(t)t^m| \leq \varepsilon$. Since $h_{\delta}(t) \in$

$C^*(A', U')$ and $t \mapsto t^m V^m = A'^m$ we deduce that $gV^m \in C^*(A', U')$ since $C^*(A', U')$ is norm closed. Thus we deduce immediately that for all $k \in \mathbb{Z}$ we have $gV^m U^k \in C^*(A', U')$, so $C_0([0, 1]) \otimes \mathcal{A}_\theta \subseteq C^*(A', U')$, hence $f_2 \in C^*(A', U')$. Hence our lemma is proven. $\square$

Theorem 5.3. *Let $\theta \in [0, 1]$. Let $\varphi : z \in \mathbb{D} \mapsto e^{2i\pi\theta} z$ where $\mathbb{D}$ is the closed unit disk in $\mathbb{C}$. Then $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ is $*$ -isomorphic to:*

$$\{f \in C([0, 1], \mathcal{A}_\theta) : f(0) \in C^*(U)\}$$

where $\mathcal{A}_\theta = C^*(U, V)$ with U, V unitaries such that $VU = e^{2i\pi\theta} UV$.

Proof. Since $\mathbb{D} \setminus \{0\}$ is fixed by φ and $C(\{0\}) \rtimes_\varphi \mathbb{Z} = C(\mathbb{T})$, we have the following easy exact sequence:

$$0 \longrightarrow C(\mathbb{D} \setminus \{0\}) \rtimes_\varphi \mathbb{Z} \longrightarrow C(\mathbb{D}) \rtimes_\varphi \mathbb{Z} \longrightarrow C(\{0\}) \rtimes_\varphi \mathbb{Z} \longrightarrow 0$$

Let $\mathfrak{B}_\theta = \{f \in C([0, 1], \mathcal{A}_\theta) : f(0) \in C^*(U)\}$. For $f \in \mathfrak{B}_\theta$ and $t \in [0, 1]$ we set $\zeta_t(f) = f(t)$. We set $A' : t \in [0, 1] \mapsto tV$ and $U' : t \in [0, 1] \mapsto U$ in $\mathfrak{B}_\theta$. Then we immediately see that $U'^* A' U' = \varphi(A')$. Hence there exists a unique $*$ -morphism ψ from $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ into $\mathfrak{B}_\theta$ such that $\psi(U_\varphi) = U'$ and $\psi(A_\varphi) = A'$. By Lemma (5.2), $\mathfrak{B}_\theta = C^*(A', U')$ so ψ is a $*$ -epimorphism onto $\mathfrak{B}_\theta$.

Now, assume $\psi(a) = 0$ for some $a \in C(\mathbb{D} \setminus \{0\}) \rtimes_\varphi \mathbb{Z}$. Let π be a nonzero irreducible $*$ -representation of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ on some Hilbert space $\mathcal{H}$. First, assume $\pi = \rho \circ \xi_r$ for some $\rho \in \widehat{\mathcal{A}_\theta}$ and some $r \in]0, 1]$. We define $\pi' = \rho \circ \zeta_t$ on $\mathfrak{B}_\theta$ and check that by construction $\pi = \pi' \circ \psi$ on the generators, hence on $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ (as $\pi, \pi' \circ \psi$ $*$ -morphisms). Therefore, $\pi(a) = 0$. Now, assume $\pi = \nu_\omega \circ \xi_0$ for some $\omega \in \mathbb{T}$. Similarly, by setting $\pi' = \nu_\omega \circ \zeta_0$ we see that $\pi(a) = \pi' \circ \psi(a) = 0$. By Proposition (5.1) we deduce that for all (irreducible) representation π of $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ we have $\pi(a) = 0$, so by definition $\|a\|_{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}} = 0$ and so $a = 0$. Therefore, ψ is a $*$ -isomorphism from $C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ onto $\mathfrak{B}_\theta$. $\square$

In particular, if $P \in C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ is a projection, then it is obviously homotopic to either 0 or 1 (for all $t, s \in [0, 1]$ set $Q(s, t) = P(st)$), so we deduce immediately that $K_0(C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}) = \mathbb{Z}$. It is also immediate that $K_1(C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}) = \mathbb{Z}$ is generated by $t \in [0, 1] \mapsto U$, as any unitary $W \in C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}$ is homotopic to a unitary of $C(\mathbb{T})$ by setting $W' : s, t \in [0, 1] \mapsto W(st)$ and $K_1(C(\mathbb{T}))$ is generated by U . Of course, these results confirm the more general facts of the previous section.

As in the case of hyperbolic automorphisms, some description of the topology of $\widehat{C(\mathbb{D}) \rtimes_\varphi \mathbb{Z}}$ is possible. To ease notations, we identify

$C(\widehat{\mathbb{D}}) \rtimes_{\varphi} \mathbb{Z}$ and $\mathfrak{B}_{\theta}$, and for all $t \in [0, 1]$ we have $\zeta_t = \xi_t$. We first define again a partition of $C(\widehat{\mathbb{D}}) \rtimes_{\varphi} \mathbb{Z}$:

Definition 5.4. *Using the notations of Proposition (5.1), we define:*

- $\text{IrrRep}_1 = \{\pi_{\omega} = \nu_{\omega} \circ \xi_0 : \omega \in \mathbb{T}\}$ and
- $\text{IrrRep}_c = \left\{ \rho \circ \xi_r : \rho \in \widehat{\mathcal{A}}_{\theta}, r \in]0, 1] \right\},$

so that we have $\text{IrrRep}_1 \cup \text{IrrRep}_c = C(\widehat{\mathbb{D}}) \rtimes_{\varphi} \mathbb{Z}$ by Proposition (5.1).

We distinguish two cases: when θ is irrational, $C(\widehat{\mathbb{D}}) \rtimes_{\varphi} \mathbb{Z}$ is not Hausdorff, yet when $\theta \in \mathbb{Q}$ then $C(\widehat{\mathbb{D}}) \rtimes_{\varphi} \mathbb{Z}$ is a compact Hausdorff space.

Proposition 5.5. *Let θ be irrational and let φ be the rotation $z \mapsto \exp(2i\pi\theta)z$. Define the map q by $\rho \circ \xi_r \in \text{IrrRep}_c \mapsto (\rho, r) \in \widehat{\mathcal{A}}_{\theta} \times]0, 1]$ and $\pi_{\omega} \in \text{IrrRep}_1 \mapsto \omega \in \mathbb{T}$ and identify IrrRep with the spectrum of $C(\widehat{\mathbb{D}}) \rtimes_{\varphi} \mathbb{Z}$ as a set.*

Then q is an homeomorphism from $C(\widehat{\mathbb{D}}) \rtimes_{\varphi} \mathbb{Z}$ onto $\mathbb{T} \cup (\widehat{\mathcal{A}}_{\theta} \times]0, 1])$ when $\mathbb{T} \cup (\widehat{\mathcal{A}}_{\theta} \times]0, 1])$ is endowed with the topology where closed sets are either of the form $\mathbb{T} \cup (\widehat{\mathcal{A}}_{\theta} \times F)$ or $Y \cup (\widehat{\mathcal{A}}_{\theta} \times X)$ where F is relatively closed in $]0, 1]$, the set X is compact in $]0, 1]$ and Y is closed in $\mathbb{T}$.

Proof. Let F be a subset of $C(\widehat{\mathbb{D}}) \rtimes_{\varphi} \mathbb{Z}$. We set

$$(5.1) \quad T = \left\{ r \in]0, 1] : \exists \rho \in \widehat{\mathcal{A}}_{\theta} \quad \rho \circ \xi_r \in F \right\}.$$

Let y be in the closure of T in $[0, 1]$. Now if $f \in C(\widehat{\mathbb{D}}) \rtimes_{\varphi} \mathbb{Z}$ such that $\pi(f) = 0$ for all $\pi \in F \cap \text{IrrRep}_c$, then $f(T) = \{0\}$, as all the representations of $\mathcal{A}_{\theta}$ are faithful (as θ irrational so $\mathcal{A}_{\theta}$ is simple), therefore $f(y) = 0$ by continuity. In particular, if 0 is in the closure of T then $\pi_{\omega}(f) = 0$ for all $\omega \in \mathbb{T}$ and thus $\text{IrrRep}_1 \subseteq F$. Moreover, since $\rho \circ \xi_y(f) = 0$ for all $f \in \bigcap_{\pi \in F} \ker \pi$ and for all $\rho \in \widehat{\mathcal{A}}_{\theta}$, if F

is closed then $\rho \circ \xi_y \in F$ for all $\rho \in \widehat{\mathcal{A}}_{\theta}$ and thus $y \in T$, so T is closed and $F \cap \text{IrrRep}_c = T \cap]0, 1] \times \widehat{\mathcal{A}}_{\theta}$. By [6, Proposition 3.2.1, p. 61], the set IrrRep_1 is closed and homeomorphic to $\mathbb{T}$ via the map $\omega \in \mathbb{T} \mapsto \pi_{\omega} \in \text{IrrRep}_1$, so if F is closed then $F \cap \text{IrrRep}_1$ is closed as well, and so is Y in $\mathbb{T}$. This proves the necessary conditions of our Proposition (note that T is compact in $]0, 1]$ if and only if it is closed in $]0, 1]$ and 0 is not in its closure in $[0, 1]$).

We now turn to the sufficient condition. Let again F be a given set in $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ and assume that T , as defined in equation (5.1), is closed in $]0, 1]$ and Y closed in $\mathbb{T}$. Now, set:

$$\mathcal{M} = \{f \in C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z} : f(T) = \{0\} \text{ and } f(0)(Y) = \{0\}\}.$$

Let $f \in \mathcal{M}$. By construction, $\pi(\mathcal{M}) = \{0\}$ for all $\pi \in F$. Conversely, let $\pi \in C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ such that $\pi(\mathcal{M}) = \{0\}$. Now, if $\pi \in \text{IrrRep}_1$, then let $\mu \in \mathbb{T}$ such that $\pi = \pi_{\mu}$. Then $\pi_{\mu}(f) = f(0)(\mu) = 0$ for all $f(0) \in C(\mathbb{T})$ such that $f(Y) = \{0\}$. Thus, $\mu \in \overline{Y} = Y$. Hence $\pi \in F$. Now, assume $\pi = \rho \circ \xi_x$ for $x \in]0, 1]$. Then $\pi(f) = 0$ by construction. If τ is the canonical tracial trace of $\mathcal{A}_{\theta}$, then it is easily seen that $\hat{\tau}(f) : t \mapsto \tau(f(t))$ is a continuous surjection from $C_0(]0, 1], \mathcal{A}_{\theta})$ onto $C_0(]0, 1])$. Now, $f \in \mathcal{M}$ if and only if $\hat{\tau}(f)(t) = 0$ for all $t \in T$. Therefore, if $\hat{\tau}(f)(x) = 0$ for all $f \in \mathcal{M}$ then $x \in \overline{T} \cap]0, 1]$. By assumption, $\{\rho \in \widehat{\mathcal{A}_{\theta}} : \rho \circ \xi_x \in F\} = \widehat{\mathcal{A}_{\theta}}$ (we assumed it was nonempty), so it follows that $\pi \in F$. Hence $F = \{\pi \in C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z} : \pi(\mathcal{M}) = \{0\}\}$, so F is closed by [6, Proposition 3.1.2, p. 60]. This concludes our proof. $\square$

Remark 5.6. *The set $\mathbb{T} \times]0, 1]$ is not in bijection with $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$, as $\widehat{\mathcal{A}_{\theta}}$ has many elements, though its topology is the coarse topology.*

Proposition 5.7. *Let $\theta \in \mathbb{Q}$. Then the spectrum of $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ is homeomorphic to $\mathfrak{S} = \{(t\alpha, \beta) : \alpha, \beta \in \mathbb{T}, t \in [0, 1]\}$ with its usual topology.*

Proof. Let $\theta = \frac{p}{q}$ with p, q relatively prime, and $\omega = \exp(2i\pi\theta)$. Let $\Xi_q = C(\mathfrak{S}, M_q(\mathbb{C}))$. For all $(\eta, \lambda) \in \mathfrak{S}$ we set:

$$U'(\eta, \lambda) = \eta \begin{bmatrix} 0 & \cdots & 0 & 1 \\ 1 & \ddots & & 0 \\ 0 & \ddots & \ddots & \vdots \\ 0 & 0 & 1 & 0 \end{bmatrix}, \quad A'(\eta, \lambda) = \begin{bmatrix} \omega & & & \\ & \omega\lambda & & \\ & & \ddots & \\ & & & \omega^{q-1}\lambda \end{bmatrix}.$$

Of course, $U', A' \in \Xi_q$ and $U'^* A' U' = \omega A' = \varphi(A')$, and $\sigma(A') = \mathbb{D}$. Therefore, there exists a $*$ -epimorphism ρ from $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ onto $C^*(A', U')$ such that $\rho(A_{\varphi}) = A'$ and $\rho(U_{\varphi}) = U'$. Using the method of the proof of Proposition (5.3), we see that ρ is also injective. Hence $C(\mathbb{D}) \rtimes_{\varphi} \mathbb{Z}$ is homeomorphic to the spectrum of $C^*(A', U')$. Note that $C^*(A', U')$ is not Ξ_q since $f \in C^*(A', U')$ implies that $f(\eta, \varphi(\lambda)) = \omega f(\eta, \lambda)$ for all $(\eta, \lambda) \in \mathfrak{S}$. Yet, it is easy to prove that $C^*(A', U') = \mathfrak{S}$. Given $\rho \in \widehat{\mathcal{A}_{\theta}}$ there exists $(\eta, \beta) \in \mathbb{T}^2$ such that $\rho = \pi_{\eta, \beta}$ where $\pi_{\eta, \beta}(U) = U'(\eta, \beta)$ and $\pi_{\eta, \beta}(V) = A'(\eta, \beta)$, so $\pi_{\eta, \beta}(A_{\varphi}(t)) = A'(\eta, t\beta)$.

Let us define $\rho_{\eta,t\beta} \in C^*(\widehat{A'}, U')$ by $\rho_{\eta,t\beta} = \pi_{\eta,\beta} \circ \zeta_t$ for all $t \in [0, 1]$ and $(\eta, \beta) \in \mathbb{T}^2$. We then see by 5.1 that $C^*(\widehat{A'}, U') = \{\rho_x : x \in \mathfrak{S}\}$ and the map $q : \rho_x \mapsto x \in \mathfrak{S}$ is an obvious bijection. Now, let F be a subset of $C^*(\widehat{A'}, U')$ and let $M = \bigcap_{\pi \in F} \ker \pi$. Let $G = q(F)$ and let $f \in M$. Then $y \in \overline{G}$ if and only if $f(y) = 0$ for all $f \in M$, and thus $M \subseteq \ker \rho_y$. Thus, if F is closed then $\rho_y \in F$ so $y \in G$ and thus G is closed; if G is closed then $y \in G$ so $\rho_y \in F$ and $F = \{\rho : \rho(M) = \{0\}\}$ so F is closed by [6, Proposition 3.1.2, p.60]. Hence q is a homeomorphism. $\square$

6. APPENDIX: SOME CALCULATIONS

We group in this appendix a few details about proofs of theorems in the main text. We include them in this version of our paper for the reader's convenience.

Proposition 6.1. *An automorphism $\varphi : z \in \mathbb{D} \mapsto e^{2i\pi\theta} \frac{z-z_0}{z-\overline{z_0}z}$ ($|z_0| < 1$, $\theta \in [0, 1)$), $\varphi \neq \text{Id}$, is hyperbolic, elliptic or parabolic if and only if, respectively, $|\sin(\pi\theta)| < |z_0|$, $|\sin(\pi\theta)| > |z_0|$ or $|\sin(\pi\theta)| = |z_0|$.*

Proof. We recall from Theorem (2.3) that the fixed point of φ are the roots of $\overline{z_0}z^2 + (e^{2i\pi\theta} - 1)z - e^{2i\pi\theta}z_0$. This polynomial is of degree 1 if and only if $z_0 = 0$, and in this case φ is a rotation (hence elliptic, unless $\theta = 0$ and then $\varphi = \text{Id}$), and we have indeed $|z_0| < |\sin(\pi\theta)|$ for $\varphi \neq \text{Id}$ and $z_0 = 0$. Otherwise, the discriminant of this trinomial is $\Delta = (e^{2i\pi\theta} - 1)^2 + 4|z_0|^2$, while the modulus of the product of the roots is $\left|e^{2i\pi\theta} \frac{z_0}{\overline{z_0}}\right| = 1$, hence either both fixed points (counting multiplicity) are on $\mathbb{T}$, or only one is in $\mathbb{D}$, and then it lies in the interior of $\mathbb{D}$. Since $e^{2i\pi\theta} - 1 = 2i \exp(i\pi\theta) (\sin(\pi\theta))$, we have $\Delta = 4 \exp(2i\pi\theta) (|z_0|^2 - \sin^2(\pi\theta))$. Hence, $\Delta = 0$ if and only if $|z_0| = |\sin(\pi\theta)|$. Now, we denote $\sqrt[2]{\Delta}$ for any square root of the complex Δ and $z_{\pm}$ the fixed points of φ , so that

$$z_{\pm} \overline{z_0} = \frac{(1 - e^{2i\pi\theta}) \pm \sqrt[2]{\Delta}}{2} = e^{i\pi\theta} \left(i \sin(\pi\theta) \pm \sqrt{|z_0|^2 - \sin^2(\pi\theta)} \right).$$

If $|z_0| > |\sin(\pi\theta)|$ then the roots $\pm \sqrt[2]{\Delta}$ of Δ are real and distinct, so

$$|z_{\pm}|^2 = |z_0|^{-2} |z_{\pm} z_0|^2 = |z_0|^{-2} ((\sin(\pi\theta))^2 + |z_0|^2 - \sin^2(\pi\theta)^2) = 1.$$

Hence, $z_{\pm} \in \mathbb{T}$ and are distinct. If $|z_0| < |\sin(\pi\theta)|$ then $|\overline{z_0}^{-1}(\lambda - 1)| = 2|z_0|^{-1} |\sin(\pi\theta)| > 2$. Yet if both fixed points of φ are in $\mathbb{T}$ then the modulus of their sum is bounded above by 2. Hence, φ has one interior fixed point in $\mathbb{D}$ (and one outside of $\mathbb{D}$). $\square$

Proposition 6.2. *Let φ be a hyperbolic automorphism of $\mathbb{D}$. Then there exists a unique $a \in (0, 1)$ such that φ is conformally conjugated to $z \mapsto \frac{z+a}{1+az}$.*

Proof. By Theorem (2.3), there exists such an $a \in (0, 1)$. To prove it is unique, we consider the following. Let $\varphi : z \mapsto \frac{z+x_0}{1+x_0z}$ and $\psi : z \mapsto \frac{z+y_0}{1+y_0z}$ for $x_0, y_0 \in (0, 1)$. Assume that there exists $\phi \in \mathbb{M}_{\mathbb{D}}$ such that $\phi \circ \psi \circ \phi^{-1} = \varphi$ and define $c \in \mathbb{D}$ such that $0 = \phi(c)$. Note that $|c| < 1$. Now, $\phi \circ \psi \circ \phi^{-1} = \varphi$ implies that $\phi(\{-1, 1\}) = \{-1, 1\}$. If $\phi(1) = 1$ then $\phi = z \mapsto \frac{z-c}{1-\bar{c}z}$ and thus ϕ commutes with φ, ψ and thus $\varphi = \psi$. If $\phi(1) = -1$, we check that $\phi = z \mapsto \lambda \frac{z-c}{1-\bar{c}z}$ for some $\lambda \in \mathbb{T}$. Since $\phi(1) = -1$ we conclude $\lambda = \frac{1-\bar{c}}{c-1}$, and since $\phi(-1) = 1$ we have $\lambda = -\frac{1+\bar{c}}{1+c}$, so $c = \bar{c}$ and thus $\lambda = -1$. Note that we then have $\phi^{-1} = \phi$. Since $\phi \circ \psi \circ \phi = \varphi$ we must have $\varphi(c) = \phi(y_0)$, hence $\frac{c+x_0}{1+x_0c} = -\frac{y_0-c}{1-cy_0}$, so $0 = x_0c^2 + cx_0y_0 - y_0 - x_0$. This is equivalent to $c \in \{-1, 1\}$ or $y_0 = -x_0$, yet both are impossible by assumption. Hence $x_0 = y_0$. $\square$

Proposition 6.3. *Any two hyperbolic automorphisms of $\mathbb{D}$ are topologically conjugated.*

Proof. We let φ and ψ be two hyperbolic automorphisms with attractive fixed point 1 and repulsive fixed point -1 , which is sufficient by Theorem (2.3).

We define for all $\infty \geq r \geq 1$ the circle Γ_r of radius r and passing by the points -1 and 1 and such that $\Gamma_r \cap \mathbb{D}$ is contained in the upper half plane. We denote by Γ_{-r} the symmetric of Γ_r with respect to the real axis. Note that $\varphi(\Gamma_{\varepsilon r}) = \psi(\Gamma_{\varepsilon r}) = \Gamma_{\varepsilon r}$ for all $\varepsilon \in \{-1, 1\}$ and $r \in [1, \infty]$. Let γ_{φ}^{ix} be the arc of Γ_{rix} from ix to $\varphi(ix)$ and let γ_{ψ}^{ix} be the arc of Γ_{rix} from ix to $\psi(ix)$, where all the arcs we consider in this proof are the ones contained in $\mathbb{D}$. Let s_{φ} and s_{ψ} be the respective length of γ_{φ}^{ix} and γ_{ψ}^{ix} when Γ_{rix} inherits its metric from the usual metric on $\mathbb{C}$. Let $\omega \in \gamma_{\varphi}^{ix}$ and let s_{ω} be the length of the arc of Γ_{rix} from ix to ω . We set $\mu_0(\omega)$ to be the unique point in Γ_{rix} and the right half plane so that the arc of Γ_r from ix to $\mu_0(\omega)$ has length $\frac{s_{\omega}}{s_{\varphi}}s_{\psi}$. Then it is easy to check that $\mu_0(\omega) \in \gamma_{\psi}^{ix}$ and $\mu_0 \circ \varphi(ix) = \psi \circ \mu_0(ix)$ by construction. Now, let $\mathcal{D}_{\varphi} = \bigcup_{x \in [-1, 1]} (\gamma_{\varphi}^{ix} \setminus \{\varphi(ix)\})$. By construction, for all $z \in \mathbb{D}$, there exists a unique $y \in \mathcal{D}_{\varphi}$ such that $\mathcal{O}_{\varphi}(z) = \mathcal{O}_{\varphi}(y)$. With obvious notations, we therefore have constructed a bijection $\mu_0 : \overline{\mathcal{D}_{\varphi}} \rightarrow \overline{\mathcal{D}_{\psi}}$ which is continuous as well (it is obvious by construction on each arc γ_{φ}^{ix} . Some easy calculation shows that μ_0 is in fact continuous on $\overline{\mathcal{D}_{\varphi}}$). Now, since our construction is symmetric in φ and ψ , we also obtain a continuous bijection $\mu'_0 : \overline{\mathcal{D}_{\psi}} \rightarrow \overline{\mathcal{D}_{\varphi}}$ which also satisfies $\mu'_0 \circ \mu_0 =$

$\mu'_0 \circ \mu_0 = \text{Id}_{\overline{\mathcal{D}_\varphi}}$ by symmetry. Hence μ_0 is a homeomorphism from $\overline{\mathcal{D}_\varphi}$ onto $\overline{\mathcal{D}_\psi}$ such that $\mu_0 \circ \varphi = \psi \circ \mu_0$ on $i[-1, 1]$.

By induction, we now extend μ_0 to a homeomorphism of $\mathbb{D}$ which conjugates φ and ψ . For all $n \in \mathbb{Z}$, we set $X_n^\varphi = \varphi^n(\overline{\mathcal{D}_\varphi})$ and $X_n^\psi = \psi^n(\overline{\mathcal{D}_\psi})$, as well as $L_n^\varphi = \varphi^n(L_0^\varphi)$. Then we assume that for some $n \in \mathbb{N}$, we have constructed the homeomorphism μ_n from X_n^φ onto X_n^ψ such that $\mu_n \circ \varphi = \psi \circ \mu_n$ on L_n^φ . We define $\mu_{n+1} = \psi \circ \mu_n \circ \varphi^{-1}$: thus μ_{n+1} is by construction an homeomorphism from X_{n+1}^φ onto X_{n+1}^ψ . Let $x \in L_{n+1}^\varphi$. We then have $\mu_{n+1}(x) = \psi \circ \mu_n \circ \varphi^{-1}(x) = \mu_n(x)$ by our induction hypothesis. Hence μ_n and μ_{n+1} agree on L_{n+1}^φ . Hence $\mu_{n+1}(\varphi(x)) = \psi(\mu_n(x)) = \psi \circ \mu_{n+1}(x)$, hence our induction is complete since we have already constructed μ_0 . The same construction yields μ_n for all $n \leq 0$ as well.

We now define the map μ by setting $\mu(x) = \mu_n(x)$ for any $x \in \mathbb{D} \setminus \{-1, 1\}$ where $n \in \mathbb{Z}$ is defined by $x \in X_n^\varphi$. Now, μ is well-defined since if $x \in X_n^\varphi \cap X_{n+1}^\varphi = L_{n+1}^\varphi$ then $\mu_{n+1}(x) = \mu_n(x)$ by the previous induction. It is now immediate that μ is an homeomorphism from $\mathbb{D} \setminus \{-1, 1\}$ onto itself. We also set $\mu(\varepsilon) = \varepsilon$ for $\varepsilon \in \{-1, 1\}$. We now check that μ is also continuous at 1 and -1 . By symmetry, it suffices to prove continuity at 1. Let $(x_n)_{n \in \mathbb{N}}$ be a sequence in $\mathbb{D}$ with limit 1. Let Ω be a neighborhood of 1 in $\mathbb{D}$. Then there exists $M \in \mathbb{N}$ such that $\cup_{j \geq M} X_j^\psi \subset \Omega$. Now there exists $N \in \mathbb{N}$ such that $x_n \in \cup_{j \geq M} X_j^\varphi$ for all $n \geq N$. Hence $\mu(x_n) \in \cup_{j \geq M} X_j^\psi \subset \Omega$ for all $n \geq N$ and thus $\lim_{n \rightarrow \infty} \mu(x_n) = 1$.

By symmetry of φ and ψ again, μ is a homeomorphism of $\mathbb{D}$. Now, let $x \in \mathbb{D} \setminus \{-1, 1\}$ and let $n \in \mathbb{Z}$ such that $x \in X_n^\varphi$. Then $\mu \circ \varphi(x) = \mu_{n+1}(\varphi(x)) = \psi(\mu_{n+1}(x)) = \psi \circ \mu(x)$, hence μ conjugates ψ and φ (by construction $\mu \circ \varphi(\varepsilon) = \psi \circ \mu(\varepsilon)$ when $\varepsilon \in \{-1, 1\}$). $\square$

REFERENCES

1. W. Arveson, *An invitation to C^* -Algebras*, GTM, vol. 39, Springer, 1976.
2. B. Blackadar, *K-theory for operator algebras*, MSRI Publications, vol. 5, 1998.
3. J. B. Conway, *A course in functional analysis*, Graduate Texts in Mathematics, vol. 96, Springer-Verlag, 1990.
4. K. R. Davidson, *C^* -algebras by examples*, Fields Institute Monographs, American Mathematical Society, 1996.
5. A. De Rango, *C^* -algebras associated with homeomorphisms of the circle*, Ph.D. thesis, University of Toronto, 2002.
6. J. Dixmier, *Les C^* -algebres et leur représentations*, Gauthier-Villars, 1969, (reprint) Editions Jacques Gabay, 1996.
7. P. Fillmore, *A user's guide to operator algebras*, Canadian Math. Soc. Series of Monographs and Adv. Texts, Wiley Interscience, 1996.

8. A. Katok and B. Hasselblatt, *Introduction to the modern theory of dynamical systems*, Cambridge University Press, 1995, 1999.
9. S. Lang, *Complex analysis*, fourth ed., GTM, vol. 103, Springer, 1999.
10. F. Latrémolière, *Approximation of the quantum tori by finite quantum tori for the quantum gromov-hausdorff distance*, Journal of Funct. Anal. **223** (2005), 365–395, math.OA/0310214.
11. S. Mac Lane, *Homology*, Classics in Mathematics, Springer, 1994.
12. G. K. Pedersen, *C^* -Algebras and their automorphism groups*, Academic Press, 1979.
13. M. Pimsner and D. V. Voiculescu, *Exact sequences for K -groups and Ext-groups of certain crossed product C^* -algebras*, Journal of Operator Theory **4** (1980), no. 1, 93–118.
14. M. A. Rieffel, *C^* -algebras associated with irrational rotations*, Pacific Journal of Mathematics **93** (1981), 415–429.
15. J. Tomiyama, *Invitation to C^* -algebras and topological dynamics*, World Scientific, 1987.
16. J. Tomiyama and S. Sylvestrov, *Topological dynamical systems of type*, Expo. Math. **20** (2002), no. 2, 117–142.
17. D. Williams, *The topology on the primitive ideal space of transformation group C^* -algebras and C^* -algebras of transformation group C^* -algebras*, Trans. Amer. Math. Soc. **266** (1981), no. 2, 335–359.
18. G. Zeller-Meier, *Produits croisés d'une C^* -algèbre par un groupe d'Automorphismes*, J. Math. pures et appl. **47** (1968), no. 2, 101–239.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO
E-mail address: `choi@math.toronto.edu`.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO
E-mail address: `frederic@math.berkeley.edu`
URL: `http://www.math.toronto.edu/frederic/`